



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07K 14/475, 14/52, 14/53, 14/535, 14/54, 14/55, 14/555, 14/56, 14/565, 14/57, C12N 15/09, 15/12, 15/18, 15/19, 15/20, 15/21, 15/22, 15/23, 15/24, 15/26, 15/27	A1	(11) International Publication Number: WO 99/03887 (43) International Publication Date: 28 January 1999 (28.01.99)
(21) International Application Number: PCT/US98/14497 (22) International Filing Date: 13 July 1998 (13.07.98) (30) Priority Data: 60/052,516 14 July 1997 (14.07.97) US (71) Applicant (for all designated States except US): BOLDER BIOTECHNOLOGY, INC. [US/US]; 678 West Willow Street, Louisville, CO 80027 (US). (72) Inventor; and (75) Inventor/Applicant (for US only): COX, George, N., III [US/US]; 678 West Willow Street, Louisville, CO 80027 (US). (74) Agent: KOIVUNIEMI, Paul, J.; 6397 Swallow Lane, Boulder, CO 80303 (US).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: DERIVATIVES OF GROWTH HORMONE AND RELATED PROTEINS		
(57) Abstract <p>The growth hormone supergene family comprises greater than 20 structurally related cytokines and growth factors. A general method is provided for creating site-specific, biologically active conjugates of these proteins. The method involves adding cysteine residues to non-essential regions of the proteins or substituting cysteine residues for non-essential amino acids in the proteins using site-directed mutagenesis and then covalently coupling a cysteine-reactive polymer or other type of cysteine-reactive moiety to the proteins via the added cysteine residue. Disclosed herein are preferred sites for adding cysteine residues or introducing cysteine substitutions into the proteins, and the proteins and protein derivatives produced thereby.</p>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon			PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

DERIVATIVES OF GROWTH HORMONE AND RELATED PROTEINS

Field of the Invention

The present invention relates to genetically engineered therapeutic proteins. More
5 specifically, the engineered proteins include growth hormone and related proteins.

Background of the Invention

The following proteins are encoded by genes of the growth hormone (GH) supergene
family (Bazan (1990); Mott and Campbell (1995); Silvennoinen and Ihle (1996)): growth
10 hormone, prolactin, placental lactogen, erythropoietin (EPO), thrombopoietin (TPO),
interleukin-2 (IL-2), IL-3, IL-4, IL-5, IL-6, IL-7, IL-9, IL-10, IL-11, IL-12 (p35 subunit), IL-
13, IL-15, oncostatin M, ciliary neurotrophic factor, leukemia inhibitory factor, alpha
interferon, beta interferon, gamma interferon, omega interferon, tau interferon, granulocyte-
colony stimulating factor (G-CSF), granulocyte-macrophage colony stimulating factor (GM-
15 CSF), macrophage colony stimulating factor (M-CSF) and cardiotrophin-1 (CT-1) ("the GH
supergene family"). It is anticipated that additional members of this gene family will be
identified in the future through gene cloning and sequencing. Members of the GH supergene
family have similar secondary and tertiary structures, despite the fact that they generally have
limited amino acid or DNA sequence identity. The shared structural features allow new
20 members of the gene family to be readily identified.

There is considerable interest on the part of patients and healthcare providers in the
development of long acting, "user-friendly" protein therapeutics. Proteins are expensive to
manufacture and, unlike conventional small molecule drugs, are not readily absorbed by the
body. Moreover, they are digested if taken orally. Therefore, natural proteins must be
25 administered by injection. After injection, most proteins are cleared rapidly from the body,
necessitating frequent, often daily, injections. Patients dislike injections, which leads to
reduced compliance and reduced drug efficacy. Some proteins, such as erythropoietin (EPO),
are effective when administered less often (three times per week for EPO) because they are
glycosylated. However, glycosylated proteins are produced using expensive mammalian cell
30 expression systems.

The length of time an injected protein remains in the body is finite and is determined
by, e.g., the protein's size and whether or not the protein contains covalent modifications such

as glycosylation. Circulating concentrations of injected proteins change constantly, often by several orders of magnitude, over a 24-hour period. Rapidly changing concentrations of protein agonists can have dramatic downstream consequences, at times under-stimulating and at other times over-stimulating target cells. Similar problems plague protein antagonists.

5 These fluctuations can lead to decreased efficacy and increased frequency of adverse side effects for protein therapeutics. The rapid clearance of recombinant proteins from the body significantly increases the amount of protein required per patient and dramatically increases the cost of treatment. The cost of human protein pharmaceuticals is expected to increase dramatically in the years ahead as new and existing drugs are approved for more disease
10 indications.

Thus, there is a need to develop protein delivery technologies that lower the costs of protein therapeutics to patients and healthcare providers. The present invention provides a solution to this problem by providing methods to prolong the circulating half-lives of protein therapeutics in the body so that the proteins do not have to be injected frequently. This
15 solution also satisfies the needs and desires of patients for protein therapeutics that are “user-friendly”, i.e., protein therapeutics that do not require frequent injections. The present invention solves these and other problems by providing biologically active, cysteine-added variants of members of the growth hormone supergene family. The invention also provides for the chemical modification of these variants with cysteine-reactive polymers or other types
20 of cysteine-reactive moieties to produce derivatives thereof and the molecules so produced.

Summary of the Invention

The present invention provides cysteine variants of members of the GH supergene family. The variants comprise a cysteine residue substituted for a nonessential amino acid of
25 the proteins. Preferably, the variants comprise a cysteine residue substituted for an amino acid selected from amino acids in the loop regions, the ends of the alpha helices, proximal to the first amphipathic helix, and distal to the final amphipathic helix or wherein the cysteine residue is added at the N-terminus or C-terminus of the proteins. Preferred sites for substitution are the N- and O-linked glycosylation sites.

30 Also provided are cysteine variants wherein the amino acid substituted for is in the A-B loop, B-C loop, the C-D loop or D-E loop of interferon/interferon-10-like members of the GH supergene family.

Also provided are cysteine variants of members of the GH supergene family wherein the cysteine residue is introduced between two amino acids in the natural protein. In particular, the cysteine residue is introduced into the loop regions, the ends of the alpha helices, proximal to the first amphipathic helix, or distal to the final amphipathic helix. Even
5 more particularly, the cysteine variant is introduced between two amino acids in an N-O-linked glycosylation site or adjacent to an amino acid in an N-linked or O-linked glycosylation site.

More particularly are provided cysteine variants wherein the loop region where the cysteine is introduced is the A-B loop, the B-C loop, the C-D loop or D-E loop of
10 interferon/interferon-10-like members of the GH supergene family.

Such cysteine substitutions or insertion mutations also can include the insertion of one or more additional amino acids at the amino-terminal or carboxy-terminal to the cysteine substitution or insertion.

Also provided are cysteine variants that are further derivatised by PEGylating the
15 cysteine variants and including the derivatised proteins produced thereby.

As set forth in the examples, specific cysteine variants of the members of the GH supergene family also are provided, including for example, variants of GH. The GH cysteine variants can have the substituted-for amino acid or inserted cysteine located at the N-terminal end of the A-B loop, the B-C loop, the C-D loop, the first three or last three amino acids in
20 the A, B, C and D helices and the amino acids proximal to helix A and distal to helix D.

More particularly, the cysteine can be substituted for the following amino acids: F1, T3, P5, E33, A34, K38, E39, K40, S43, Q46, N47, P48, Q49, T50, S51, S55, T60, A98, N99, S100, G104, A105, S106, E129, D130, G131, S132, P133, T135, G136, Q137, K140, Q141, T142, S144, K145, D147, T148, N149, S150, H151, N152, D153, S184, E186, G187, S188,
25 and G190.

Other examples of cysteine variants according to the invention include erythropoietin variants. Erythropoietin variants include those wherein the substituted for amino acid is located in the A-B loop, the B-C loop, the C-D loop, the amino acids proximal to helix A and distal to helix D and the N- or C-terminus. Even more specifically, the EPO cysteine variants
30 include molecules wherein the amino acids indicated below have a cysteine substituted therefor: serine-126, N24, I25, T26, N38, I39, T40, N83, S84, A1, P2, P3, R4, D8, S9, T27, G28, A30, E31, H32, S34, N36, D43, T44, K45, N47, A50, K52, E55, G57, Q58, G77, Q78,

A79, Q86, W88, E89, T107, R110, A111, G113, A114, Q115, K116, E117, A118, S120, P121, P122, D123, A124, A125, A127, A128, T132, K154, T157, G158, E159, A160, T163, G164, D165, R166 and S85.

The members of the GH supergene family include growth hormone, prolactin,
5 placental lactogen, erythropoietin, thrombopoietin, interleukin-2, interleukin-3, interleukin-4, interleukin-5, interleukin-6, interleukin-7, interleukin-9, interleukin-10, interleukin-11, interleukin-12 (p35 subunit), interleukin-13, interleukin-15, oncostatin M, ciliary neurotrophic factor, leukemia inhibitory factor, alpha interferon, beta interferon, gamma interferon, omega interferon, tau interferon, granulocyte-colony stimulating factor,
10 granulocyte-macrophage colony stimulating factor, macrophage colony stimulating factor, cardiotrophin-1 and other proteins identified and classified as members of the family. The proteins can be derived from any animal species including human, companion animals and farm animals.

Other variations and modifications to the invention will be obvious to those skilled in
15 the art based on the specification and the “rules” set forth herein. All of these are considered as part of the invention.

Detailed Description of the Invention

The present invention relates to cysteine variants and, among other things, the site-
20 specific conjugation of such proteins with polyethylene glycol (PEG) or other such moieties. PEG is a non-antigenic, inert polymer that significantly prolongs the length of time a protein circulates in the body. This allows the protein to be effective for a longer period of time. Covalent modification of proteins with PEG has proven to be a useful method to extend the circulating half-lives of proteins in the body (Abuchowski et al., 1984; Hershfield, 1987;
25 Meyers et al., 1991). Covalent attachment of PEG to a protein increases the protein’s effective size and reduces its rate of clearance rate from the body. PEGs are commercially available in several sizes, allowing the circulating half-lives of PEG-modified proteins to be tailored for individual indications through use of different size PEGs. Other benefits of PEG modification include an increase in protein solubility, an increase in *in vivo* protein stability
30 and a decrease in protein immunogenicity (Katre et al., 1987; Katre, 1990).

The preferred method for PEGylating proteins is to covalently attach PEG to cysteine residues using cysteine-reactive PEGs. A number of highly specific, cysteine-reactive PEGs

with different reactive groups (e.g., maleimide, vinylsulfone) and different size PEGs (2-20 kDa) are commercially available (e.g., from Shearwater, Polymers, Inc., Huntsville, AL). At neutral pH, these PEG reagents selectively attach to “free” cysteine residues, i.e., cysteine residues not involved in disulfide bonds. The conjugates are hydrolytically stable. Use of

5 cysteine-reactive PEGs allows the development of homogeneous PEG-protein conjugates of defined structure.

Considerable progress has been made in recent years in determining the structures of commercially important protein therapeutics and understanding how they interact with their protein targets, e.g., cell-surface receptors, proteases, etc. This structural information can be

10 used to design PEG-protein conjugates using cysteine-reactive PEGs. Cysteine residues in most proteins participate in disulfide bonds and are not available for PEGylation using cysteine-reactive PEGs. Through in vitro mutagenesis using recombinant DNA techniques, additional cysteine residues can be introduced anywhere into the protein. The added cysteines can be introduced at the beginning of the protein, at the end of the protein, between two

15 amino acids in the protein sequence or, preferably, substituted for an existing amino acid in the protein sequence. The newly added “free” cysteines can serve as sites for the specific attachment of a PEG molecule using cysteine-reactive PEGs. The added cysteine must be exposed on the protein’s surface and accessible for PEGylation for this method to be successful. If the site used to introduce an added cysteine site is non-essential for biological

20 activity, then the PEGylated protein will display essentially wild type (normal) in vitro bioactivity. The major technical challenge in PEGylating proteins with cysteine-reactive PEGs is the identification of surface exposed, non-essential regions in the target protein where cysteine residues can be added or substituted for existing amino acids without loss of bioactivity.

25 Cysteine-added variants of a few human proteins and PEG-polymer conjugates of these proteins have been described. US patent 5,206,344 describes cysteine-added variants of IL-2. These cysteine-added variants are located within the first 20 amino acids from the amino terminus of the mature IL-2 polypeptide chain. The preferred cysteine variant is at position 3 of the mature polypeptide chain, which corresponds to a threonine residue that is

30 O-glycosylated in the naturally occurring protein. Substitution of cysteine for threonine at position 3 yields an IL-2 variant that can be PEGylated with a cysteine-reactive PEG and retain full in vitro bioactivity (Goodson and Katre, 1990). In contrast, natural IL-2 PEGylated

with lysine-reactive PEGs displays reduced in vitro bioactivity (Goodson and Katre, 1990). The effects of cysteine substitutions at other positions in IL-2 were not reported.

US patent 5,166,322 teaches cysteine-added variants of IL-3. These variants are located within the first 14 amino acids from the N-terminus of the mature protein sequence.

5 The patent teaches expression of the proteins in bacteria and covalent modification of the proteins with cysteine-reactive PEGs. No information is provided as to whether the cysteine-added variants and PEG-conjugates of IL-3 are biologically active. Cysteine-added variants at other positions in the polypeptide chain were not reported.

World patent application WO9412219 and PCT application US95/06540 teach
10 cysteine-added variants of insulin-like growth factor-I (IGF-I). IGF-I has a very different structure from GH and is not a member of the GH supergene family (Mott and Campbell, 1995). Cysteine substitutions at many positions in the IGF-I protein are described. Only certain of the cysteine-added variants are biologically active. The preferred site for the cysteine added variant is at amino acid position 69 in the mature protein chain. Cysteine
15 substitutions at positions near the N-terminus of the protein (residues 1-3) yielded IGF-I variants with reduced biological activities and improper disulfide bonds.

World patent application WO9422466 teaches two cysteine-added variants of insulin-like growth factor (IGF) binding protein-1, which has a very different structure than GH and is not a member of the GH supergene family. The two cysteine-added IGF binding protein-1
20 variants disclosed are located at positions 98 and 101 in the mature protein chain and correspond to serine residues that are phosphorylated in the naturally-occurring protein.

US patent application 07/822296 teaches cysteine added variants of tumor necrosis factor binding protein, which is a soluble, truncated form of the tumor necrosis factor cellular receptor. Tumor necrosis factor binding protein has a very different structure than GH and is
25 not a member of the GH supergene family.

IGF-I, IGF binding protein-1 and tumor necrosis factor binding protein have secondary and tertiary structures that are very different from GH and the proteins are not members of the GH supergene family. Because of this, it is difficult to use the information gained from studies of IGF-I, IGF binding protein-1 and tumor necrosis factor binding protein
30 to create cysteine-added variants of members of the GH supergene family. The studies with IL-2 and IL-3 were carried out before the structures of IL-2 and IL-3 were known (McKay 1992; Bazan, 1992) and before it was known that these proteins are members of the GH

supergene family. Previous experiments aimed at identifying preferred sites for adding cysteine residues to IL-2 and IL-3 were largely empirical and were performed prior to experiments indicating that members of the GH supergene family possessed similar secondary and tertiary structures.

5 Based on the structural information now available for members of the GH supergene family, the present invention provides “rules” for determining a priori which regions and amino acid residues in members of the GH supergene family can be used to introduce or substitute cysteine residues without significant loss of biological activity. In contrast to the naturally occurring proteins, these cysteine-added variants of members of the GH supergene
10 family will possess novel properties such as the ability to be covalently modified at defined sites within the polypeptide chain with cysteine-reactive polymers or other types of cysteine-reactive moieties. The covalently modified proteins will be biologically active.

GH is the best-studied member of the GH supergene family. GH is a 22 kDa protein secreted by the pituitary gland. GH stimulates metabolism of bone, cartilage and muscle and
15 is the body’s primary hormone for stimulating somatic growth during childhood. Recombinant human GH (rhGH) is used to treat short stature resulting from GH inadequacy and renal failure in children. GH is not glycosylated and can be produced in a fully active form in bacteria. The protein has a short in vivo half-life and must be administered by daily subcutaneous injection for maximum effectiveness (MacGillivray et al., 1996). Recombinant
20 human GH (rhGH) was approved recently for treating cachexia in AIDS patients and is under study for treating cachexia associated with other diseases.

The sequence of human GH is well known (see, e.g., Martial et al. 1979; Goeddel et al. 1979 which are incorporated herein by reference; SEQ ID NO:1). GH is closely related in sequence to prolactin and placental lactogen and these three proteins were considered
25 originally to comprise a small gene family. The primary sequence of GH is highly conserved among animal species (Abdel-Meguid et al., 1987), consistent with the protein’s broad species cross-reactivity. The three dimensional folding pattern of porcine GH has been solved by X-ray crystallography (Abdel-Meguid et al., 1987). The protein has a compact globular structure, comprising four amphipathic alpha helical bundles joined by loops.
30 Human GH has a similar structure (de Vos et al., 1992). The four alpha helical regions are termed A-D beginning from the N-terminus of the protein. The loop regions are referred to by the helical regions they join, e.g., the A-B loop joins helical bundles A and B. The A-B

and C-D loops are long, whereas the B-C loop is short. GH contains four cysteine residues, all of which participate in disulfide bonds. The disulfide assignments are cysteine53 joined to cysteine165 and cysteine182 joined to cysteine189.

The crystal structure of GH bound to its receptor revealed that GH has two receptor
5 binding sites and binds two receptor molecules (Cunningham et al., 1991; de Vos et al., 1992). The two receptor binding sites are referred to as site I and site II. Site I encompasses the Carboxy (C)-terminal end of helix D and parts of helix A and the A-B loop, whereas site II encompasses the Amino (N)-terminal region of helix A and a portion of helix C. Binding of GH to its receptor occurs sequentially, with site I always binding first. Site II then engages
10 a second GH receptor, resulting in receptor dimerization and activation of the intracellular signaling pathways that lead to cellular responses to GH. A GH mutein in which site II has been mutated (a glycine to arginine mutation at amino acid 120) is able to bind a single GH receptor, but is unable to dimerize GH receptors; this mutein acts as a GH antagonist *in vitro*, presumably by occupying GH receptor sites without activating intracellular signaling
15 pathways (Fuh et al., 1992).

The roles of particular regions and amino acids in GH receptor binding and intracellular signaling also have been studied using techniques such as mutagenesis, monoclonal antibodies and proteolytic digestion. The first mutagenesis experiments entailed replacing entire domains of GH with similar regions of the closely related protein, prolactin
20 (Cunningham et al., 1989). One finding was that replacement of the B-C loop of GH with that of prolactin did not affect binding of the hybrid GH protein to a soluble form of the human GH receptor, implying that the B-C loop was non-essential for receptor binding. Alanine scanning mutagenesis (replacement of individual amino acids with alanine) identified 14 amino acids that are critical for GH bioactivity (Cunningham and Wells, 1989). These
25 amino acids are located in the helices A, B, C, and D and the A-B loop and correspond to sites I and II identified from the structural studies. Two lysine residues at amino acid positions 41 and 172, K41 and K172, were determined to be critical components of the site I receptor binding site, which explains the decrease in bioactivity observed when K172 is acetylated (Teh and Chapman, 1988). Modification of K168 also significantly reduced GH
30 receptor binding and bioactivity (de la Llosa et al., 1985; Martal et al., 1985; Teh and Chapman, 1988). Regions of GH responsible for binding the GH receptor have also been studied using monoclonal antibodies (Cunningham et al., 1989). A series of eight

monoclonal antibodies was generated to human GH and analyzed for the ability to neutralize GH activity and prevent binding of GH to its recombinant soluble receptor. The latter studies allowed the putative binding site for each monoclonal antibody to be localized within the GH three-dimensional structure. Of interest was that monoclonal antibodies 1 and 8 were unable to displace GH from binding its receptor. The binding sites for these monoclonal antibodies were localized to the B-C loop (monoclonal number 1) and the N-terminal end of the A-B loop (monoclonal number 8). No monoclonals were studied that bound the C-D loop specifically. The monoclonal antibody studies suggest that the B-C loop and N-terminal end of the A-B loop are non-essential for receptor binding. Finally, limited cleavage of GH with trypsin was found to produce a two chain derivative that retained full activity (Mills et al., 1980; Li, 1982). Mapping studies indicated that trypsin cleaved and/or deleted amino acids between positions 134 and 149, which corresponds to the C-D loop. These studies suggest the C-D loop is not involved in receptor binding or GH bioactivity.

Structures of a number of cytokines, including G-CSF (Hill et al., 1993), GM-CSF (Diederichs et al., 1991; Walter et al., 1992), IL-2 (Bazan, 1992; McKay, 1992), IL-4 (Redfield et al., 1991; Powers et al., 1992), and IL-5 (Milburn et al., 1993) have been determined by X-ray diffraction and NMR studies and show striking conservation with the GH structure, despite a lack of significant primary sequence homology. EPO is considered to be a member of this family based upon modeling and mutagenesis studies (Boissel et al., 1993; Wen et al., 1994). A large number of additional cytokines and growth factors including ciliary neurotrophic factor (CNTF), leukemia inhibitory factor (LIF), thrombopoietin (TPO), oncostatin M, macrophage colony stimulating factor (M-CSF), IL-3, IL-6, IL-7, IL-9, IL-12, IL-13, IL-15, and alpha, beta, omega, tau and gamma interferon belong to this family (reviewed in Mott and Campbell, 1995; Silvennoinen and Ihle 1996). All of the above cytokines and growth factors are now considered to comprise one large gene family, of which GH is the prototype.

In addition to sharing similar secondary and tertiary structures, members of this family share the property that they must oligomerize cell surface receptors to activate intracellular signaling pathways. Some GH family members, e.g., GH and EPO, bind a single type of receptor and cause it to form homodimers. Other family members, e.g., IL-2, IL-4, and IL-6, bind more than one type of receptor and cause the receptors to form heterodimers or higher order aggregates (Davis et al., 1993; Paonessa et al., 1995; Mott and Campbell, 1995).

Mutagenesis studies have shown that, like GH, these other cytokines and growth factors contain multiple receptor binding sites, typically two, and bind their cognate receptors sequentially (Mott and Campbell, 1995; Matthews et al., 1996). Like GH, the primary receptor binding sites for these other family members occur primarily in the four alpha helices and the A-B loop (reviewed in Mott and Campbell, 1995). The specific amino acids in the helical bundles that participate in receptor binding differ amongst the family members (Mott and Campbell, 1995). Most of the cell surface receptors that interact with members of the GH supergene family are structurally related and comprise a second large multi-gene family (Bazan, 1990; Mott and Campbell, 1995; Silvennoinen and Ihle 1996).

A general conclusion reached from mutational studies of various members of the GH supergene family is that the loops joining the alpha helices generally tend to not be involved in receptor binding. In particular the short B-C loop appears to be non-essential for receptor binding in most, if not all, family members. For this reason, the B-C loop is a preferred region for introducing cysteine substitutions in members of the GH supergene family. The A-B loop, the B-C loop, the C-D loop (and D-E loop of interferon/ IL-10-like members of the GH superfamily) also are preferred sites for introducing cysteine mutations. Amino acids proximal to helix A and distal to the final helix also tend not to be involved in receptor binding and also are preferred sites for introducing cysteine substitutions. Certain members of the GH family, e.g., EPO, IL-2, IL-3, IL-4, IL-6, G-CSF, GM-CSF, TPO, IL-10, IL-12 p35, IL-13, IL-15 and beta-interferon contain N-linked and O-linked sugars. The glycosylation sites in the proteins occur almost exclusively in the loop regions and not in the alpha helical bundles. Because the loop regions generally are not involved in receptor binding and because they are sites for the covalent attachment of sugar groups, they are preferred sites for introducing cysteine substitutions into the proteins. Amino acids that comprise the N- and O-linked glycosylation sites in the proteins are preferred sites for cysteine substitutions because these amino acids are surface-exposed, the natural protein can tolerate bulky sugar groups attached to the proteins at these sites and the glycosylation sites tend to be located away from the receptor binding sites.

Many additional members of the GH gene family are likely to be discovered in the future. New members of the GH supergene family can be identified through computer-aided secondary and tertiary structure analyses of the predicted protein sequences. Members of the GH supergene family will possess four or five amphipathic helices joined by non-helical

amino acids (the loop regions). The proteins may contain a hydrophobic signal sequence at their N-terminus to promote secretion from the cell. Such later discovered members of the GH supergen family also are included within this invention.

The present invention provides “rules” for creating biologically active cysteine-added
5 variants of members of the GH supergene family. These “rules” can be applied to any existing or future member of the GH supergene family. The cysteine-added variants will possess novel properties not shared by the naturally occurring proteins. Most importantly, the cysteine added variants will possess the property that they can be covalently modified with cysteine-reactive polymers or other types of cysteine-reactive moieties to generate biologically active
10 proteins with improved properties such as increased in vivo half-life, increased solubility and improved in vivo efficacy.

Specifically, the present invention provides biologically active cysteine variants of members of the GH supergene family by substituting cysteine residues for non-essential amino acids in the proteins. Preferably, the cysteine residues are substituted for amino acids
15 that comprise the loop regions, for amino acids near the ends of the alpha helices and for amino acids proximal to the first amphipathic helix or distal to the final amphipathic helix of these proteins. Other preferred sites for adding cysteine residues are at the N-terminus or C-terminus of the proteins. Cysteine residues also can be introduced between two amino acids in the disclosed regions of the polypeptide chain. The present invention teaches that N- and
20 O-linked glycosylation sites in the proteins are preferred sites for introducing cysteine substitutions either by substitution for amino acids that make up the sites or, in the case of N-linked sites, introduction of cysteines therein. The glycosylation sites can be serine or threonine residues that are O-glycosylated or asparagine residues that are N-glycosylated. N-linked glycosylation sites have the general structure asparagine-X-serine or threonine (N-X-S/T), where X can be any amino acid. The asparagine residue, the amino acid in the X
25 position and the serine/threonine residue of the N-linked glycosylation site are preferred sites for creating biologically active cysteine-added variants of these proteins. Amino acids immediately surrounding or adjacent to the O-linked and N-linked glycosylation sites (within about 10 residues on either side of the glycosylation site) are preferred sites for introducing
30 cysteine-substitutions.

More generally, certain of the “rules” for identifying preferred sites for creating biologically active cysteine-added protein variants can be applied to any protein, not just

proteins that are members of the GH supergene family. Specifically, preferred sites for creating biologically active cysteine variants of proteins (other than IL-2) are O-linked glycosylation sites. Amino acids immediately surrounding the O-linked glycosylation site (within about 10 residues on either side of the glycosylation site) also are preferred sites. N-linked glycosylation sites, and the amino acid residues immediately adjacent on either side of the glycosylation site (within about 10 residues of the N-X-S/T site) also are preferred sites for creating cysteine added protein variants. Amino acids that can be replaced with cysteine without significant loss of biological activity also are preferred sites for creating cysteine-added protein variants. Such non-essential amino acids can be identified by performing cysteine-scanning mutagenesis on the target protein and measuring effects on biological activity. Cysteine-scanning mutagenesis entails adding or substituting cysteine residues for individual amino acids in the polypeptide chain and determining the effect of the cysteine substitution on biological activity. Cysteine scanning mutagenesis is similar to alanine-scanning mutagenesis (Cunningham et al., 1992), except that target amino acids are individually replaced with cysteine rather than alanine residues.

Application of the “rules” to create cysteine-added variants and conjugates of protein antagonists also is contemplated. Excess production of cytokines and growth factors has been implicated in the pathology of many inflammatory conditions such as rheumatoid arthritis, asthma, allergies and wound scarring. Excess production of GH has been implicated as a cause of acromegaly. Certain growth factors and cytokines, e.g., GH and IL-6, have been implicated in proliferation of particular cancers. Many of the growth factors and cytokines implicated in inflammation and cancer are members of the GH supergene family. There is considerable interest in developing protein antagonists of these molecules to treat these diseases. One strategy involves engineering the cytokines and growth factors so that they can bind to, but not oligomerize receptors. This is accomplished by mutagenizing the second receptor binding site (site II) on the molecules. The resulting muteins are able to bind and occupy receptor sites but are incapable of activating intracellular signaling pathways. This strategy has been successfully applied to GH to make a GH antagonist (Cunningham et al., 1992). Similar strategies are being pursued to develop antagonists of other members of the GH supergene family such as IL-2 (Zurawski et al., 1990; Zurawski and Zurawski, 1992), IL-4 (Kruse et al., 1992), IL-5 (Tavernier et al., 1995), GM-CSF (Hercus et al., 1994) and EPO (Matthews et al., 1996). Since the preferred sites for adding cysteine residues to members of

the GH supergene family described here lie outside of the receptor binding sites in these proteins, and thus removed from any sites used to create protein antagonists, the cysteine-added variants described herein could be used to generate long-acting versions of protein antagonists. As an example, Cunningham et al. (1992) developed an in vitro GH antagonist
5 by mutating a glycine residue (amino acid 120) to an arginine. This glycine residue is a critical component of the second receptor binding site in GH; when it is replaced with arginine, GH cannot dimerize receptors. The glycine to arginine mutation at position 120 can be introduced into DNA sequences encoding the cysteine-added variants of GH contemplated herein to create a cysteine-added GH antagonist that can be conjugated with cysteine-reactive
10 PEGs or other types of cysteine-reactive moieties. Similarly, amino acid changes in other proteins that turn the proteins from agonists to antagonists could be incorporated into DNA sequences encoding cysteine-added protein variants described herein. Considerable effort is being spent to identify amino acid changes that convert protein agonists to antagonists. Hercus et al.(1994) reported that substituting arginine or lysine for glutamic acid at position
15 21 in the mature GM-CSF protein converts GM-CSF from an agonist to an antagonist. Tavernier et al.(1995) reported that substituting glutamine for glutamic acid at position 13 of mature IL-5 creates an IL-5 antagonist.

Experimental strategies similar to those described above can be used to create cysteine-added variants (both agonists and antagonists) of members of the GH supergene
20 family derived from various animals. This is possible because the primary amino acid sequences and structures of cytokines and growth factors are largely conserved between human and animal species. For this reason, the “rules” disclosed herein for creating biologically active cysteine-added variants of members of the GH supergene family will be useful for creating biologically active cysteine-added variants of members of the GH
25 supergene family of companion animals (e.g., dogs, cats, horses) and commercial animal (e.g., cow, sheep, pig) species. Conjugation of these cysteine-added variants with cysteine-reactive PEGs will create long-acting versions of these proteins that will benefit the companion animal and commercial farm animal markets.

Proteins that are members of the GH supergene family (hematopoietic cytokines) are
30 provided in Silvennoimem and Ihle (1996). Silvennoimem and Ihle (1996) also provide information about the structure and expression of these proteins. DNA sequences, encoded amino acids and in vitro and in vivo bioassays for the proteins described herein are described

in Aggarwal and Gutterman (1992; 1996), Aggarwal (1998), and Silvennoimem and Ihle (1996). Bioassays for the proteins also are provided in catalogues of various commercial suppliers of these proteins such as R&D Systems, Inc. and Endogen, Inc.

The following examples are provided to demonstrate how these “rules” can be used to create cysteine-added variants of GH, erythropoietin, alpha interferon, beta interferon, G-CSF, GM-CSF and other members of the GH supergene family. The examples are not intended to be limiting, but only exemplary of specific embodiments of the invention.

Example 1

Cysteine-added variants of GH

This example discloses certain amino acids in GH that are non-essential for biological activity and which, when mutated to cysteine residues, will not alter the normal disulfide binding pattern and overall conformation of the molecule. These amino acids are located at the N-terminal end of the A-B loop (amino acids 34-52 of the mature protein sequence; SEQ ID NO: 1; Martial et al 1979; Goeddel et al 1979), the B-C loop (amino acids 97-105 in the mature protein sequence), and the C-D loop (amino acids 130-153 in the mature protein sequence). Also identified as preferred sites for introducing cysteine residues are the first three or last three amino acids in the A, B, C and D helices and the amino acids proximal to helix A and distal to helix D.

DNA sequences encoding wild type GH can be amplified using the polymerase chain reaction technique from commercially available single-stranded cDNA prepared from human pituitaries (ClonTech, San Diego, CA) or assembled using overlapping oligonucleotides. Specific mutations can be introduced into the GH sequence using a variety of procedures such as phage techniques (Kunkel et al 1987), PCR mutagenesis techniques (Innis et al 1990; White 1993) mutagenesis kits such as those sold by Stratagene (“Quick-Change Mutagenesis” kit, San Diego, CA) or Promega (Gene Editor Kit, Madison WI).

Cysteine substitutions can be introduced into any of the amino acids comprising the B-C loop, C-D loop and N-terminal end of the A-B loop or into the first three amino acids of the alpha helical regions that adjoin these regions. or in the region proximal to helix A or distal to helix D. Preferred sites for introduction of cysteine residues are: F1, T3, P5, E33, A34, K38, E39, Q40, S43, Q46, N47, P48, Q49, T50, S51, S55, T60, A98, N99, S100, G104, A105, S106, E129, D130, G131, S132, P133, T135, G136, Q137, K140, Q141, T142, S144,

K145, D147, T148, N149, S150, H151, N152, D153, S184, E186, G187, S188, and G190.

Cysteine residues also can be introduced at the beginning of the mature protein, i.e., proximal to the F1 amino acid, or following the last amino acid in the mature protein, i.e., following F191. If desirable, two or more such mutations can be readily combined in the same protein either by in vitro DNA recombination of cloned mutant genes and/or sequential construction of individual desired mutations.

1. Cloning the Gene for human Growth Hormone (GH)

The human GH gene was amplified from human pituitary single-stranded cDNA (commercially available from CLONTECH, Inc., Palo Alto, CA) using the polymerase chain reaction (PCR) technique and primers BB1 and BB2. The sequence of BB1 is 5'-GGGGGTCGACCATATGTTCCCAACCATTCCTTATCCAG-3' (SEQ ID NO: 24). The sequence of BB2 is 5'-GGGGGATCCTCACTAGAAGCCACAGCTGCCCTC-3' (SEQ ID NO: 25). Primer BB1 was designed to encode an initiator methionine preceding the first amino acid of mature GH, phenylalanine, and Sall and NdeI sites for cloning purposes. The reverse primer, BB2, contains a BamHI site for cloning purposes. The PCR 100 microliter reactions contained 20 pmoles of each oligonucleotide primer, 1X PCR buffer (Perkin-Elmer buffer containing MgCl₂), 200 micromolar concentration of each of the four nucleotides dA, dC, dG and dT, 2 ng of single-stranded cDNA, 2.5 units of Taq polymerase (Perkin-Elmer) and 2.5 units of Pfu polymerase (Stratagene, Inc). The PCR reaction conditions were 96°C for 3 minutes, 35 cycles of (95°C, 1 minute; 63°C for 30 seconds; 72°C for 1 minute), followed by 10 minutes at 72°C. The thermocycler employed was the Amplitron II Thermal Cycler (Thermolyne). The approximate 600 bp PCR product was digested with Sall and BamHI, gel purified and cloned into similarly digested plasmid pUC19 (commercially available from New England BioLabs, Beverly, MA). The ligation mixture was transformed into E. coli strain DH5alpha and transformants selected on LB plates containing ampicillin. Several colonies were grown overnight in LB media and plasmid DNA isolated using miniplasmid DNA isolation kits purchased from Qiagen, Inc (Valencia, CA). Clone LB6 was determined to have the correct DNA sequence.

For expression in E. coli, clone LB6 was digested with NdeI and EcoRI, the approximate 600 bp fragment gel-purified, and cloned into plasmid pCYB1 (commercially available from New England BioLabs, Beverly, MA) that had been digested with the same enzymes and phosphatased. The ligation mixture was transformed into E. coli DH5alpha and

transformants selected on LB ampicillin plates. Plasmid DNA was isolated from several transformants and screened by digestion with NdeI and EcoRI. A correct clone was identified and named pCYB1: wtGH (pBBT120). This plasmid was transformed into E. coli strains JM109 or W3110 (available from New England BioLabs and the American Type Culture
5 Collection).

2. Construction of STII-GH

Wild type GH clone LB6 (pUC19: wild type GH) was used as the template to construct a GH clone containing the E. coli STII signal sequence (Picken et al. 1983).

Because of its length, the STII sequence was added in two sequential PCR reactions. The
10 first reaction used forward primer BB12 and reverse primer BB10. BB10 has the sequence: 5'CGCGGATCCGATTAGAATCCACAGCTCCCCTC 3' (SEQ ID NO: 28).

BB12 has the sequence:

5'ATCTATGTTCGTTTTCTCTATCGCTACCAACGCTTACGCATTCCCAACCATTCCC
TTATCCAG-3' (SEQ ID NO: 30).

15 The PCR reactions were as described for amplifying wild type GH except that approximately 4 ng of plasmid LB6 was used as the template rather than single-stranded cDNA and the PCR conditions were 96⁰C for 3 minutes, 30 cycles of (95⁰C for 1 minute; 63⁰C for 30 seconds; 72⁰C for 1 minute) followed by 72⁰C for 10 minutes. The approximate 630 bp PCR product was gel-purified using the Qiaex II Gel Extraction Kit (Qiagen, Inc),
20 diluted 50-fold in water and 2 microliters used as template for the second PCR reaction. The second PCR reaction used reverse primer BB10 and forward primer BB11. BB11 has the sequence:

5'CCCCCTCTAGACATATGAAGAAGAACATCGCATTCCTGCTGGCATCTATGTTCG
TTTTCTCTATCG-3' (SEQ ID NO: 29).

25 Primer BB11 contains XbaI and NdeI sites for cloning purposes. PCR conditions were as described for the first reaction. The approximate 660 bp PCR product was digested with XbaI and BamHI, gel-purified and cloned into similarly cut plasmid pCDNA3.1(+) (Invitrogen, Inc. Carlsbad, CA). Clone pCDNA3.1(+):stII-GH(5C) or "5C" was determined to have the correct DNA sequence.

30 Clone "5C" was cleaved with NdeI and BamHI and cloned into similarly cut pBBT108 (a derivative of pUC19 which lacks a Pst I site, this plasmid is described below). A clone with the correct insert was identified following digestion with these enzymes. This

clone, designated pBBT111, was digested with NdeI and Sall, the 660 bp fragment containing the stII-GH fusion gene, was gel-purified and cloned into the plasmid expression vector pCYB1 (New England BioLabs) that had been digested with the same enzymes and phosphatased. A recombinant plasmid containing the stII-GH insertion was identified by
5 restriction endonuclease digestions. One such isolate was chosen for further studies and was designated pBBT114. This plasmid was transformed into E. coli strains JM109 or W3110 (available from New England BioLabs and the American Type Culture Collection).

3. Construction of ompA-GH

Wild type GH clone LB6 (pUC19: wild type GH) was used as the template to
10 construct a GH clone containing the E. coli ompA signal sequence (Movva et al 1980).

Because of its length, the ompA sequence was added in two sequential PCR reactions. The first reaction used forward primer BB7:

5'GCAGTGGCACTGGCTGGTTTCGCTACCGTAGCGCAGGCCTTCCCAACCATTCCC
TTATCCAG 3' (SEQ ID NO: 31),

15 and reverse primer BB10:

5' CGCGGATCCGATTAGAATCCACAGCTCCCCTC 3' (SEQ ID NO: 28).

The PCR reactions were as described for amplifying wild type GH except that approximately 4 ng of plasmid LB6 was used as the template rather than single-stranded cDNA and the PCR conditions were 96⁰C for 3 minutes, 30 cycles of (95⁰C for 1 minute;
20 63⁰C for 30 seconds; 72⁰C for 1 minute) followed by 72⁰C for 10 minutes. The approximate 630 bp PCR product was gel-purified using the Qiaex II Gel Extraction Kit (Qiagen, Inc), diluted 50-fold in water and 2 microliters used as template for the second PCR reaction. The second PCR reaction used reverse primer BB10 and forward PrimerBB6:

5'CCCCGTCGACACATATGAAGAAGACAGCTATCGCGATTGCAGTGGCACTGGCT
25 GGTTTC 3' (SEQ ID NO: 32).

PCR conditions were as described for the first reaction. The approximate 660 bp PCR product was gel-purified, digested with Sal I and Bam H1 and cloned into pUC19 (New England BioLabs) which was cut with Sal I and Bam H1 or pCDNA3.1(+)
(Invitrogen) which had been cut by Xho I and Bam H1 (Sal I and Xho I produce compatible
30 single-stranded overhangs). When several clones were sequenced, it was discovered that all pUC19 clones (8/8) contained errors in the region of the ompA sequence. Only one pCDNA3.1(+) clone was sequenced and it contained a sequence ambiguity in the ompA

region. In order to generate a correct ompA-GH fusion gene segments of two sequenced clones which contained different errors separated by a convenient restriction site were recombined and cloned into the pUC19-derivative that lacks the Pst I site (see pBBT108 described below). The resulting plasmid, termed pBBT112, carries the ompA-GH fusion gene
5 cloned as an Nde I - Bam H1 fragment into these same sites in pBBT108. This plasmid is designated pBBT112 and is used in PCR-based, site-specific mutagenesis of GH as described below.

4. Construction of Pst- pUC19

To facilitate mutagenesis of the cloned GH gene for construction of selected cysteine
10 substitution and insertion mutations a derivative of the plasmid pUC19 (New England BioLabs) lacking a Pst I site was constructed as follows. pUC19 plasmid DNA was digested with Pst I and subsequently treated at 75 deg. C with PFU DNA Polymerase (Stratagene) using the vendor-supplied reaction buffer supplemented with 200uM dNTPs. Under these conditions the polymerase will digest the 3' single-stranded overhang created by Pst I
15 digestion but will not digest into the double-stranded region. The net result will be the deletion of the 4 single-stranded bases which comprise the middle four bases of the Pst I recognition site. The resulting molecule has double-stranded, i.e., "blunt", ends. Following these enzymatic reactions, the linear monomer was gel-purified using the Qiaex II Gel Extraction Kit (Qiagen, Inc). This purified DNA was treated with T4 DNA Ligase (New
20 England BioLabs) according to the vendor protocols, digested with Pst I, and used to transform E coli DH5alpha. Transformants were picked and analyzed by restriction digestion with Pst I and Bam H1. One of the transformants which was not cleaved by Pst I but was cleaved at the nearby Bam H1 site was picked and designated pBBT108.

5. Construction of GH muteins

25 GH muteins were generally constructed using site-directed PCR-based mutagenesis as described in PCR Protocols: Current Methods and Applications edited by B. A. White, 1993 Humana Press, Inc., Totowa, NJ and PCR Protocols: A Guide to Methods and Applications edited by Innis, M. A. et al 1990 Academic Press Inc San Diego, CA . Typically PCR primer oligonucleotides are designed to incorporate nucleotide changes to the coding sequence of
30 GH that result in substitution of a cysteine residue for an amino acid at a specific position within the protein. Such mutagenic oligonucleotide primers can also be designed to incorporate an additional cysteine residue at the carboxy terminus or amino terminus of the

coding sequence of GH. In this latter case one or more additional amino acid residues could also be incorporated at the amino terminal and/or carboxy terminal to the added cysteine residue if that were desirable. Moreover, oligonucleotides can be designed to incorporate cysteine residues as insertion mutations at specific positions within the GH coding sequence if that were desirable. Again, one or more additional amino acids could be inserted along with the cysteine residue and these amino acids could be positioned at the amino terminal and/or carboxy terminal to the cysteine residue.

The cysteine substitution mutation T135C was constructed as follows. The mutagenic reverse oligonucleotide BB28:

5'CTGCTTGAAGATCTGCCCACACCGGGGGCTGCCATC3' (SEQ ID NO: 33)

was designed to change the codon ACT for threonine at amino acid residue 135 to a TGT codon encoding cysteine and to span the nearby Bgl II site. This oligonucleotide was used in PCR along with the forward oligonucleotide BB34

5'GTAGCGCAGGCCTTCCCAACCATT3' (SEQ ID NO: 34) which anneals to the junction region of the ompA-GH fusion gene and is not mutagenic. The PCR was performed in a 50ul reaction in 1X PCR buffer (Perkin-Elmer buffer containing 1.5 mM MgCl₂), 200 micromolar concentration of each of the four nucleotides dA, dC, dG and dT, with each oligonucleotide primer present at 0.5 μM, 5 pg of pBBT112 (described above) as template and 1.25 units of AmpliTac DNA Polymerase (Perkin-Elmer) and 0.125 units of PFU DNA Polymerase

(Stratagene). Reactions were performed in a Robocycler Gradient 96 thermal cycler (Stratagene). The program used entailed: 95 deg C for 3 minutes followed by 25 cycles of 95 deg C for 60 seconds, 45 deg C or 50 deg C or 55 deg C for 75 seconds, 72 deg C for 60 seconds followed by a hold at 6 deg C. The PCR reactions were analyzed by agarose gel electrophoresis to identify annealing temperatures that gave significant product of the expected size; ~430 bp. The 45-deg C reaction was "cleaned up" using the QIAquick PCR Purification Kit (Qiagen), digested with Bgl II and Pst I. The resulting 278 bp Bgl II- Pst I fragment, which includes the putative T135C mutation, was gel-purified and ligated into pBBT111 the pUC19 derivative carrying the stII-GH fusion gene (described above) which had been digested with Bgl II and Pst I and gel-purified. Transformants from this ligation were initially screened by digestion with Bgl II and Pst I and subsequently one clone was sequenced to confirm the presence of the T135C mutation and the absence of any additional

mutations that could potentially be introduced by the PCR reaction or by the synthetic oligonucleotides. The sequenced clone was found to have the correct sequence.

The substitution mutation S132C was constructed using the protocol described above for T135C with the following differences: mutagenic reverse oligonucleotide BB29

5'CTGCTTGAAGATCTGCCCAGTCCGGGGGCAGCCATCTTC3' (SEQ ID NO: 35) was used instead of BB28 and the PCR reaction with annealing temperature of 50 deg C was used for cloning. One of two clones sequenced was found to have the correct sequence.

The substitution mutation T148C was constructed using an analogous protocol but employing a different cloning strategy. The mutagenic forward oligonucleotide BB30

5'GGGCAGATCTTCAAGCAGACCTACAGCAAGTTCGACTGCAACTCACACAAC3' (SEQ ID NO: 36) was used in PCR with the non-mutagenic reverse primer BB33

5'CGCGGTACCCGGGATCCGATTAGAATCCACAGCT3' (SEQ ID NO: 37) which anneals to the most 3' end of the GH coding sequence and spans the Bam H1 site immediately downstream. PCR was performed as described above with the exception that the

annealing temperatures used were 46, 51 and 56 deg C. Following PCR and gel analysis as described above the 46 and 51 deg C reactions were pooled for cloning. These were digested with Bam H1 and Bgl II, gel-purified and cloned into pBBT111 which had been digested with Bam H1 and Bgl II, treated with Calf intestinal Alkaline Phosphatase (Promega) according to the vendor protocols, and gel-purified. Transformants from this ligation were analyzed by

digestion with Bam H1 and Bgl II to identify clones in which the 188 bp Bam H1- Bgl II mutagenic PCR fragment was cloned in the proper orientation. Because Bam H1 and Bgl II generate compatible ends, this cloning step is not orientation specific. Five of six clones tested were shown to be correctly oriented. One of these was sequenced and was shown to contain the desired T148C mutation. The sequence of the remainder of the 188 bp Bam H1-

Bgl II mutagenic PCR fragment in this clone was confirmed as correct.

The construction of the substitution mutation S144C was identical to the construction of T148C with the following exceptions. Mutagenic forward oligonucleotide BB31

5'GGGCAGATCTTCAAGCAGACCTACTGCAAGTTCGAC3' (SEQ ID NO: 38) was used instead of BB30. Two of six clones tested were shown to be correctly oriented. One of these was sequenced and was shown to contain the desired S144C mutation. The sequence of the

remainder of the 188 bp Bam H1- Bgl II mutagenic PCR fragment in this clone was confirmed as correct.

A mutation was also constructed that added a cysteine residue to the natural carboxy terminus of GH. The construction of this mutation, termed stp192C, was similar to that of T148C, but employed different oligonucleotide primers. The reverse mutagenic oligonucleotide BB32

5'CGCGGTACCGGATCCTTAGCAGAAGCCACAGCTGCCCTCCAC3' (SEQ ID NO: 39) which inserts a TGC codon for cysteine between the codon for the carboxy terminal phe residue of GH and the TAA translational stop codon and spans the nearby Bam H1 site was used along with BB34 5'GTAGCGCAGGCCTTCCCAACCATT3' (SEQ ID NO: 40) which is described above. Following PCR and gel analysis as described above, the 46 deg C reaction was used for cloning. Three of six clones tested were shown to be correctly oriented. One of these was sequenced and was shown to contain the desired stp192C mutation. The sequence of the remainder of the 188 bp Bam H1- Bgl II mutagenic PCR fragment in this clone was confirmed as correct.

Analogous PCR mutagenesis procedures can be used to generate other cysteine mutations. The choice of sequences for mutagenic oligonucleotides will be dictated by the position where the desired cysteine residue is to be placed and the propinquity of useful restriction endonuclease sites. Generally, it is desirable to place the mutation, i.e., the mismatched segment near the middle of the oligonucleotide to enhance the annealing of the oligonucleotide to the template. Appropriate annealing temperatures for any oligonucleotide can be determined empirically. It is also desirable for the mutagenic oligonucleotide to span a unique restriction site so that the PCR product can be cleaved to generate a fragment that can be readily cloned into a suitable vector, e.g., one that can be used to express the mutein or provides convenient restriction sites for excising the mutated gene and readily cloning it into such an expression vector. Sometimes mutation sites and restriction sites are separated by distances that are greater than that which is desirable for synthesis of synthetic oligonucleotides: it is generally desirable to keep such oligonucleotides under 80 bases in length and lengths of 30-40 bases are more preferable.

In instances where this is not possible, genes targeted for mutagenesis could be re-engineered or re-synthesized to incorporate restriction sites at appropriate positions.

Alternatively, variations of PCR mutagenesis protocols employed above, such as the so-called

“Megaprimer Method” (Barik, S., pp. 277-286 in Methods in Molecular Biology, Vol. 15: PCR Protocols: Current Methods and Applications edited by B. A. White, 1993, Humana Press, Inc., Totowa, NJ) or “Gene Splicing by Overlap Extension” (Horton, R. M., pp. 251-261, in Methods in Molecular Biology, Vol. 15: PCR Protocols: Current Methods and Applications, edited by B. A. White, 1993, Humana Press, Inc., Totowa, NJ) can also be employed to construct such mutations.

6. Expression of GH in pCYB1

To express GH in *E. coli*, pBBT120 (GH gene with no leader sequence cloned into the *tac* expression vector pCYB1) and pBBT114 (GH gene with *stII* leader sequence cloned into the *tac* expression vector pCYB1) were transformed into *E. coli* strains JM109 and W3110. The parental vector pCYB1 was also transformed into JM109 and W3110.

These strains were given the following designations:

BOB119:	JM109 (pCYB1)
BOB130:	W3110 (pCYB1)
BOB129:	JM109 (pBBT120)
BOB133:	W3110 (pBBT120)
BOB121:	JM109 (pBBT114)
BOB132:	W3110 (pBBT114)

For expression, strains were grown overnight at 37 ° C in Luria Broth (LB) (Sambrook, et al 1989) containing 100 µg/ml ampicillin. These saturated overnight cultures were diluted to ~ 0.03 OD at A₆₀₀ in LB containing 100 µg/ml ampicillin and incubated at 37 ° C in shake flasks in rotary shaker, typically at 250-300 rpm. ODs were monitored and IPTG was added to a final concentration of 0.5 mM when culture ODs reached ~0.25 -0.5, typically between 0.3 and 0.4. Cultures were sampled typically at 1, 3, 5 and ~16 h post-induction. The “~16 h” time points represented overnight incubation of the cultures and exact times varied from ~15-20 h. Samples of induced and uninduced cultures were pelleted by centrifugation, resuspended in 1X sample buffer (50mM Tris-HCl (pH 6.8), 2% sodium lauryl sulfate, 10% glycerol, 0.1% bromphenol blue) with the addition of 1% β-mercaptoethanol when desirable. Samples were boiled for ~10 minutes or heated to 95 ° C for ~ 10 minutes. Samples were cooled to room temperature before being loaded onto SDS polyacrylamide gels or were stored at -20° C if not run immediately. Samples were run on precast 15% polyacrylamide “Ready Gels” (Bio-Rad, Hercules CA) using a Ready Gel Cell electrophoresis apparatus (Bio-Rad)

according to the vendor protocols. Typically gels were run at 200 volts for ~35-45 minutes. Gels were stained with Coomassie Blue or were analyzed by Western Blot following electro-blotting. Coomassie staining of whole cell lysates from strains BOB 129, BOB 133, BOB 121 and BOB 132 showed a band of ~22 kD that co-migrated with purified recombinant human GH standard purchased from Research Diagnostics Inc (Flanders, NJ). That band was most prominent in induced cultures following overnight induction. However, a band was also observed at that molecular weight in uninduced cultures of these same strains and could also be observed with and without induction in the BOB 119 and BOB 130 control strains that carried the expression vector pCYB1 lacking the GH gene. To clarify this observation, Western Blot analyses were performed on whole cell lysates of induced cultures of strains BOB119, BOB130, BOB129, BOB133, BOB121, and BOB132. Western blots were performed with polyclonal rabbit anti-human GH antiserum purchased from United States Biological; catalogue # G 9000-11 (Swampscott, MA) This primary antibody was used at a 1:5000 dilution and its binding was detected with goat anti-rabbit IgG Fc conjugated to alkaline phosphatase, (product # 31341) purchased from Pierce (Rockford, IL). This secondary antibody was used at a 1:10,000 dilution. Alkaline phosphatase activity was detected using the ImmunoPure® Fast Red TR/AS-MX Substrate Kit (Pierce, Rockford IL) according to the vendor protocols. The Western Blots clearly demonstrated presence of GH in lysates of induced cultures of BOB129, BOB133, BOB121, and BOB132 at both 3 and 16 h post-induction. In the induced culture of control strains, BOB119 and BOB130, no GH was detected by Western blot at 3 or 16 h post-induction time points.

In these preliminary experiments, the highest yields of GH were obtained from BOB132 W3110(pBBT114) in which the GH gene is fused downstream of the stII secretion signal sequence. This strain was tested further to determine if the GH protein was secreted to the periplasm as would be expected. An induced culture of BOB132 was prepared as described above and subjected to osmotic shock according to the procedure of Koshland and Botstein (Cell 20 (1980) pp. 749-760). This procedure ruptures the outer membrane and releases the contents of the periplasm into the surrounding medium. Subsequent centrifugation separates the periplasmic contents, present in the supernatant from the remainder of the cell-associated components. In this experiment, the bulk of the GH synthesized by BOB132 was found to be localized to the periplasm. This result is consistent with the finding that the bulk of the total GH is also indistinguishable in size from the

purified GH standard, which indicated that the still signal sequence had been removed. This is indicative of secretion. A larger scale (500 ml) culture of BOB132 was also induced, cultured overnight and subjected to osmotic shock according to the procedure described by Hsiung et al, 1986 (Bio/Technology 4, pp. 991-995). Gel analysis again demonstrated that the bulk of the GH produced was soluble, periplasmic, and indistinguishable in size from the GH standard. This material could also be quantitatively bound to, and eluted from, a Q-Sephadex column using conditions very similar to those described for recombinant human GH by Becker and Hsiung, 1986 (FEBS Lett 204 pp145-150).

7. Cloning the human GH receptor

The human GH receptor was cloned by PCR using forward primer BB3 and reverse primer BB4. BB3 has the sequence:

5'-CCCCGATCCGCCACCATGGATCTCTGGCAGCTGCTGTT-3' (SEQ ID NO: 26).

BB4 has the sequence:

5'CCCCGTCGACTCTAGAGCTATTAAATACGTAGCTCTTGGG-3' (SEQ ID NO: 27).

The template was single-stranded cDNA prepared from human liver (commercially available from CLONTECH Laboratories). Primers BB3 and BB4 contain BamHI and SalI restriction sites, respectively, for cloning purposes. The 100 µl PCR reactions contained 2.5 ng of the single-stranded cDNA and 20 picomoles of each primer in 1X PCR buffer (Perkin-Elmer buffer containing MgCl₂), 200 micromolar concentration of each of the four nucleotides dA, dC, dG and dT, 2.5 units of Taq polymerase (Perkin-Elmer) and 2.5 units of Pfu polymerase (Stratagene, Inc). The PCR reaction conditions were: 96°C for 3 minutes, 35 cycles of (95°C, 1 minute; 58°C for 30 seconds; 72°C for 2 minutes), followed by 10 minutes at 72°C. The thermocycler employed was the AmpliTron II Thermal Cycler (Thermolyne). The approximate 1.9 kb PCR product was digested with BamHI and SalI and ligated with similarly cut plasmid pUC19 (New England BioLabs). However, none of the transformants obtained from this ligation reaction contained the 1.9 kb PCR fragment. Leung et al (Nature 1987 330 pp537-543) also failed to obtain full-length cDNA clones of the human GH receptor in pUC19. Subsequently the PCR fragment was cloned into a low copy number vector, pACYC184 (New England BioLabs) at the BamHI and SalI sites in this vector. Such clones were obtained at reasonable frequencies but E coli strains carrying the cloned PCR fragment grew poorly, forming tiny and heterogeneous looking colonies, in the presence of chloramphenicol, which is used to select for maintenance of pACYC184.

The PCR fragment was simultaneously cloned into pCDNA3.1 (+) (Invitrogen). The approximate 1.9 kb PCR product was digested with BamHI and SalI and ligated into the BamHI and XhoI cloning sites of pCDNA3.1 (+). Only infrequent transformants from this ligation contained the cloned GH receptor cDNA and all of those were found to contain deletions of segments of the receptor coding sequence. One of these clones was sequenced and found to contain a deletion of 135 bp within the GH receptor coding sequence: the sequence of the rest of the gene was in agreement with that reported by Leung et al (1987).

8. Cloning the rabbit GH receptor

The rabbit GH receptor was cloned by PCR using forward primer BB3 (described above) and reverse primer BB36. BB36 has the sequence 5'CCCCGTCGACTCTAGAGCCATTAGATACAAAGCTCTTGGG3' (SEQ ID NO: 41) and contains XbaI and SalI restriction sites for cloning purposes. Rabbit liver poly(A)⁺ mRNA was purchased from CLONTECH, Inc. and used as the substrate in first strand synthesis of single-stranded cDNA to produce template for PCR amplification. First strand synthesis of single-stranded cDNA was accomplished using a 1st Strand cDNA Synthesis Kit for RT-PCR (AMV) kit from Boehringer Mannheim Corp (Indianapolis, IN) according to the vendor protocols. Parallel first strand cDNA syntheses were performed using random hexamers or BB36 as the primer. Subsequent PCR reactions with the products of the first strand syntheses as templates, were carried out with primers BB3 and BB36 according to the 1st Strand cDNA Synthesis Kit for RT-PCR (AMV) kit protocol and using 2.5 units of Amplitac DNA Polymerase (Perkin-Elmer) and 0.625 units of Pfu DNA Polymerase (Stratagene). The PCR reaction conditions were 96°C for 3 minutes, 35 cycles of (95°C, 1 minute; 58°C for 30 seconds; 72°C for 2 minutes), followed by 10 minutes at 72°C. The thermocycler employed was the AmpliTron II Thermal Cycler (Thermolyne). The expected ~ 1.9 kb PCR product was observed in PCR reactions using random hexamer-primed or BB36 primed cDNA as template. The random hexamer-primed cDNA was used in subsequent cloning experiments. It was digested with BamHI and XbaI and run out over a 1.2% agarose gel. This digest generates two fragments (~ 365 bp and ~ 1600 bp) because the rabbit GH receptor gene contains an internal BamHI site. Both fragments were gel-purified. Initially the ~1600 bp BamHI - XbaI fragment was cloned into pCDNA3.1(+) which had been digested with these same two enzymes. These clones were readily obtained at reasonable frequencies and showed no evidence of deletions as determined by restriction digests and

subsequent sequencing. To generate a full length clone, one of the plasmids containing the 1600 bp Bam H1 - Xba I fragment (pCDNA3.1(+): rab-ghr-2A) was digested with Bam H1, treated with Calf Intestinal Alkaline Phosphatase (Promega) according to the vendor protocols, gel-purified and ligated with the gel purified ~365 bp Bam H1 fragment that contains the 5' portion of the rabbit GH receptor gene. Transformants from this ligation were picked and analyzed by restriction digestion and PCR to confirm the presence of the ~365 bp fragment and to determine its orientation relative to the distal segment of the rabbit GH receptor gene. Three out of four clones analyzed were found to contain the ~365 bp fragment cloned in the correct orientation for reconstitution of the rabbit GH receptor gene. The lack of complications in the cloning in *E coli* of the rabbit gene, in contrast to the human gene, is consistent with the results of Leung et al (1987) who also readily obtained full length cDNA clones for the rabbit GH receptor gene but were unable to clone a full length cDNA of the human gene in *E coli*. The rabbit GH receptor can be employed in assays with human GH as a ligand as it has been shown that the human GH binds the rabbit receptor with high affinity (Leung et al 1987). Plasmids containing the cloned rabbit GH receptor should be sequenced to identify a rabbit GH receptor cDNA with the correct sequence before use.

9. Construction of a human/rabbit Chimeric GH Receptor Gene

As an alternative to the rabbit receptor, a chimeric receptor could be constructed which combines the extracellular domain of the human receptor with the transmembrane and cytoplasmic domains of the rabbit receptor. Such a chimeric receptor could be constructed by recombining the human and rabbit genes at the unique Nco I site that is present in each (Leung et al 1987). Such a recombinant, containing the human gene segment located 5' to, or "upstream" of the Nco I site and the rabbit gene segment 3' to, or "downstream" of the Nco I site would encode a chimeric receptor of precisely the desired type, having the extracellular domain of the human receptor with the transmembrane and cytoplasmic domains of the rabbit receptor. This would allow analysis of the interaction of GH, GH muteins, and PEGylated GH muteins with the natural receptor binding site but could avoid the necessity of cloning the full length human GH receptor in *E coli*.

The GH muteins can be expressed in a variety of expression systems such as bacteria, yeast or insect cells. Vectors for expressing GH muteins in these systems are available commercially from a number of suppliers such as Novagen, Inc. (pET15b for expression in *E. coli*), New England Biolabs (pC4B1 for expression in *E. coli*) Invitrogen (pVL1392,

pVL1393, and pMELBAC for expression in insect cells using Baculovirus vectors, Pichia vectors for expression in yeast cells, and pCDNA3 for expression in mammalian cells). GH has been successfully produced in E. coli as a cytoplasmic protein and as a secreted, periplasmic, protein using the E. coli OmpA or STII signal sequences to promote secretion of the protein into the periplasmic space (Chang et al., 1987; Hsiung et al., 1986). It is preferable that the GH muteins are expressed as secreted proteins so that they do not contain an N-terminal methionine residue, which is not present in the natural human protein. For expression in E. coli, DNA sequences encoding GH or GH muteins can be cloned into E. coli expression vectors such as pET15b that uses the strong T7 promoter or pCYB1 that uses the TAC promoter. Adding IPTG (isopropylthiogalactopyranoside, available from Sigma Chemical Company) to the growth media, can induce expression of the protein. Recombinant GH will be secreted into the periplasmic space from which it can be released by, and subsequently purified, following osmotic shock (Becker and Hsiung, 1986). The protein can be purified further using other chromatographic methods such as ion-exchange, hydrophobic interaction, size-exclusion and reversed phase chromatography, all of which are well known to those of skill in the art (e.g., see Becker and Hsiung, 1986). Protein concentrations can be determined using commercially available protein assay kits such as those sold by BioRad Laboratories (Richmond, CA). If the GH proteins are insoluble when expressed in E. coli they can be refolded using procedures well known to those skilled in the art (see Cox et al., 1994 and World patent applications WO9422466 and WO9412219).

Alternatively, the proteins can be expressed in insect cells as secreted proteins. The expression plasmid can be modified to contain the GH signal sequence to promote secretion of the protein into the medium. The cDNAs can be cloned into commercially available vectors, e.g., pVL1392 from Invitrogen, Inc., and used to infect insect cells. The GH and GH muteins can be purified from conditioned media using conventional chromatography procedures. Antibodies to rhGH can be used in conjunction with Western blots to localize fractions containing the GH proteins during chromatography. Alternatively, fractions containing GH can be identified using ELISA assays.

The cysteine-added GH variants also can be expressed as intracellular or secreted proteins in eukaryotic cells such as yeast, insect cells or mammalian cells. Vectors for expressing the proteins and methods for performing such experiments are described in catalogues from various commercial supply companies such as Invitrogen, Inc., Stratagene,

Inc. and ClonTech, Inc. The GH and GH muteins can be purified using conventional chromatography procedures.

Biological activity of the GH muteins can be measured using a cell line that proliferates in response to GH. Fuh et al. (1992) created a GH-responsive cell line by stably transforming a myeloid leukemia cell line, FDC-P1, with a chimeric receptor comprising the extracellular domain of the rabbit GH receptor fused to the mouse G-CSF receptor. This cell line proliferates in response to GH with a half maximal effective concentration (EC_{50}) of 20 picomolar. A similar cell line can be constructed using the published sequences of these receptors and standard molecular biology techniques (Fuh et al., 1992). Alternatively, the extracellular domain of the human GH receptor can be fused to the mouse G-CSF receptor using the published sequences of these receptors and standard molecular biology techniques. Transformed cells expressing the chimeric receptor can be identified by flow cytometry using labeled GH, by the ability of transformed cells to bind radiolabeled GH, or by the ability of transformed cells to proliferate in response to added GH. Purified GH and GH muteins can be tested in cell proliferation assays using cells expressing the chimeric receptor to measure specific activities of the proteins. Cells can be plated in 96-well dishes with various concentrations of GH or GH muteins. After 18 h, cells are treated for 4 hours with 3H -thymidine and harvested for determination of incorporated radioactivity. The EC_{50} can be determined for each mutein. Assays should be performed at least three times for each mutein using triplicate wells for each data point. GH muteins displaying similar optimal levels of stimulation and EC_{50} values comparable to or greater than wild type GH are preferable.

GH muteins that retain in vitro activity can be PEGylated using a cysteine-reactive 8 kDa PEG-maleimide (or PEG-vinylsulfone) commercially available from Shearwater, Inc. Generally, methods for PEGylating the proteins with these reagents will be similar to those described in world patent applications WO 9412219 and WO 9422466 and PCT application US95/06540, with minor modifications. The recombinant proteins must be partially reduced with dithiothreitol (DTT) in order to achieve optimal PEGylation of the free cysteine. Although the free cysteine is not involved in a disulfide bond, it is relatively unreactive to cysteine-reactive PEGs unless this partial reduction step is performed. The amount of DTT required to partially reduce each mutein can be determined empirically, using a range of DTT concentrations. Typically, a 5-10 fold molar excess of DTT for 30 min at room temperature is sufficient. Partial reduction can be detected by a slight shift in the elution profile of the

protein from a reversed-phase column. Care must be taken not to over-reduce the protein and expose additional cysteine residues. Over-reduction can be detected by reversed phase-HPLC (the protein will have a retention time similar to the fully reduced and denatured protein) and by the appearance of GH molecules containing two PEGs (detectable by a molecular weight change on SDS-PAGE). Wild type GH can serve as a control since it should not PEGylate under similar conditions. Excess DTT can be removed by size exclusion chromatography using spin columns. The partially reduced protein can be reacted with various concentrations of PEG-maleimide (PEG: protein molar ratios of 1:1, 5:1, 10:1 and 50:1) to determine the optimum ratio of the two reagents. PEGylation of the protein can be monitored by a molecular weight shift using sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE). The lowest amount of PEG that gives significant quantities of mono-pegylated product without giving di-pegylated product will be considered optimum (80% conversion to mono-pegylated product is considered good). Generally, mono-PEGylated protein can be purified from non-PEGylated protein and unreacted PEG by size-exclusion or ion exchange chromatography. The purified PEGylated protein can be tested in the cell proliferation assay described above to determine its specific activity.

The above experiments will allow identification of amino acids in the B-C loop, C-D loop or N-terminal end of the A-B loop in GH that can be changed to cysteine residues, PEGylated and retain in vitro biological activity. These muteins can be tested in animal disease models well known in the art.

Experiments can be performed to confirm that the PEG molecule is attached to the protein at the proper site. This can be accomplished by proteolytic digestion of the protein, purification of the PEG peptide (which will have a large molecular weight) by size exclusion, ion exchange or reversed phase chromatography, followed by amino acid sequencing or mass spectroscopy. The PEG-coupled amino acid will appear as a blank in the amino acid sequencing run.

The pharmacokinetic properties of the PEG-GH proteins can be determined as follows, or as described in world patent application WO9422466. Pairs of rats or mice can receive an intravenous bolus injection of the test proteins. Circulating levels of the proteins are measured over the course of 24 h by removing a small sample of blood from the animals at desired time points. Circulating levels of the test proteins can be quantitated using ELISA assays. Additional experiments can be performed using the subcutaneous route to administer

the proteins. Similar experiments should be performed with the non-PEGylated protein to serve as a control. These experiments will reveal whether attachment of a PEG reagent to the protein alters its pharmacokinetic properties. Covalent modification of the protein with PEG should increase the protein's circulating half-life relative to the unPEGylated protein. Larger
5 PEG molecules and/or attachment of multiple PEG molecules should lengthen the circulating half-life longer than smaller PEG molecules.

PEG-GH proteins can be tested in rodent models of growth hormone deficiency (Cox et al., 1994) and cachexia (Tomas et al., 1992; Read et al., 1992) to determine optimum dosing schedules and demonstrate efficacy. These studies can explore different size PEG
10 molecules, e.g., 8 and 20 kDa, and dosing schedules to determine the optimum PEG size and dosing schedule. It is expected that the larger PEG molecule will increase the circulating half-life greater than the smaller PEG molecule and will require less frequent dosing.

However, large proteins potentially may have reduced volumes of distribution *in vivo*; thus, it is possible a 20 kDa PEG attached to GH will limit bioavailability, reducing its efficacy.

15 Rodent models will allow determination of whether this is the case. Once the optimum dosing schedules and PEG sizes are determined, the efficacy of PEG-GH to GH can be compared in the animal models. While all PEG-GH proteins having GH activity are included in the invention, the preferred PEG-GH proteins are those that enhance growth equal or superior to GH, but which can be given less frequently. PEG-GH should be more efficacious
20 than GH when both are administered using the less frequent dosing schedules.

One GH deficiency model that can be used is a hypophysectomized rat. GH stimulates body weight gain and bone and cartilage growth in this model (Cox et al., 1994). Hypophysectomized rats can be purchased from Charles River. Rats can be injected with GH, PEG-GH or placebo and weight gain measured daily over a 10-14 day period. At time of
25 sacrifice, tibial epiphysis width can be determined as a measure of bone growth.

Experimental methods for performing these studies are described in Cox et al. (1994).

The efficacy of PEG-GH in rodent cachexia models can be tested in a similar manner. Daily administration of dexamethasone, via osmotic pumps or subcutaneous injection, can be used to induce weight loss (Tomas et al., 1992; Read et al., 1992; PCT patent application
30 US95/06540).

Example 2

Cysteine-added variants of Erythropoietin

This example relates to cysteine-added variants of erythropoietin (EPO). EPO is the hormone primarily responsible for stimulating erythropoiesis or red blood cell formation.

5 EPO acts on immature red blood cell precursors to stimulate their further proliferation and differentiation into mature red blood cells. A commercial pharmaceutical version is available from Amgen, Inc. Human EPO is a 35-39 kDa glycoprotein secreted by the adult kidney. The mature human protein contains 166 amino acids and is heavily glycosylated. The sequence of human EPO (SEQ ID NO: 2) is shown in Lin et al 1985 and Jacobs et al. 1985,
10 which are incorporated herein by reference. The primary sequence of EPO is highly conserved among species (greater than 80% identity; Wen et al., 1994). Sugar groups account for greater than 40% of the protein's mass. Human EPO contains three N-linked glycosylation sites and one O-linked glycosylation site. The N-linked glycosylation sites are conserved in different species whereas the O-linked glycosylation site is not. The extensive
15 glycosylation of EPO has prevented the protein's crystallization, so the X-ray structure of the protein is not known. Human EPO contains four cysteine residues. The disulfide assignments are Cys7 to Cys161 and Cys29 to Cys33. Cys33 is not conserved in mouse EPO, suggesting that the Cys29-Cys33 disulfide bond is not critical to mouse EPO's structure or function. This conclusion also seems to hold for human EPO (Boissel et al., 1993).

20 The amino acid sequence of EPO is consistent with the protein being a member of the GH supergene family and mutational studies support this view of EPO's structure (Boissel et al., 1993; Wen et al., 1994). A model of the three dimensional structure of EPO, modeled after the GH structure has been proposed (Boissel et al., 1993; Wen et al., 1994). Amino acids in EPO important for receptor binding have been identified through mutagenesis
25 experiments and reside primarily in the N-terminal half of presumptive helix A and the C-terminal half of presumptive helix D (Boissel et al., 1993; Wen et al., 1994; Matthews et al., 1996). Only a single cell surface receptor for EPO has been identified (D'Andrea et al., 1989). It is believed that EPO dimerizes its receptor in much the same way that GH dimerizes its receptor (Matthew's et al., 1996).

30 Human EPO contains three sites for N-linked glycosylation (asparagine- 24,- 38 and - 83) and one site for O-linked glycosylation (serine-126). The N-linked glycosylation sites are located in the A-B and B-C loops and the O-glycosylation site is located in the C-D loop.

The N-linked glycosylation sites are conserved among species whereas the O-linked glycosylation site is absent in rodent EPO (Wen et al., 1993). A non-O-linked glycosylated human variant containing methionine at position 126 has been described (US patent 4,703,008). The N-linked sugar groups are heavily branched and contain terminal sialic acid residues (Sasaki et al., 1987; Takeuchi et al., 1988). N-38 and N-83 contain the most highly branched oligosaccharides (Sasaki et al., 1988).

The terminal sialic residues on EPO are critical for the protein's in vivo function because removal of these residues by digestion eliminates in vivo activity (Fukada et al., 1989; Spivak and Hogans, 1989). Loss of activity correlates with faster clearance of the asialated protein from the body. The circulating half-life of the asialated protein in rats is less than ten minutes, in contrast to that of the sialated protein, which is approximately 2 hr (Fukada et al., 1989; Spivak and Hogans, 1989). Thus, in vivo activity of EPO directly correlates with its circulating half-life.

The role of N-linked sugars in EPO's biological activities has been better defined by mutating, individually and in combination, the three asparagine residues comprising the N-linked glycosylation sites. EPO muteins in which only a single N-linked glycosylation site was mutated, i.e., N24Q, N38Q and N83Q, were secreted from mammalian cells as efficiently as wild type EPO, indicating that N-linked glycosylation at all three sites is not required for protein secretion (N24Q indicates that the asparagine at position 24 is mutated to glutamine). In contrast, EPO muteins in which two or more N-linked glycosylation sites were mutated were secreted less efficiently than wild type EPO from mammalian cells (Yamaguchi et al., 1991; Delorme et al., 1992). Mutagenesis studies found that each of the single N-linked glycosylation site muteins had in vitro biological activities equal to or greater than wild type EPO. Thus, it was concluded that none of the N-linked glycosylation sites is essential for secretion or in vitro biological activity of EPO. In fact, removal of one of the glycosylation sites seemed to improve biological activity (Yamaguchi et al., 1991).

The in vivo biological activity of N-linked glycosylation muteins was studied by two groups. Yamaguchi et al. (1991) concluded that the N24Q and N83Q muteins had in vivo activities greater than wild type EPO, which correlated with their increased in vitro activities. These authors found that the N38Q mutein had decreased in vivo activity, about 60% of wild type EPO. N38 is the most heavily branched of the three N-linked glycosylation sites (Sasaki et al., 1988). Delorme et al. (1992) reported that mutating any of the N-linked glycosylation

sites reduced in vivo biological activity by about 50%. Muteins in which two or more glycosylation sites were mutated had decreased in vivo activities in both studies.

The above studies indicate that some N-linked glycosylation is required for in vitro and in vivo activity of EPO. Individually, however, none of the three glycosylation sites is absolutely essential for activity. The N-linked sugars increase the apparent molecular weight of EPO and prolong its circulating half-life, which correlates with bioactivity. Natural EPO and EPO manufactured in mammalian cells have complex N-linked sugars containing galactose and terminal sialic acid residues. The galactose residues are recognized by specific receptors on hepatocytes and promote rapid clearance of EPO from the body unless the galactose residues are masked by the terminal sialic acid residues.

Mutagenesis studies concluded that O-linked glycosylation is not required for in vitro or in vivo function of EPO (Delorme et al., 1992). This is in keeping with the observation that rodent EPO is not O-glycosylated and with the existence of a naturally occurring human EPO variant in which serine-126 is replaced by methionine, with a corresponding lack of O-linked glycosylation. Mutagenesis of serine-126 revealed that certain amino acid changes at this site (to valine, histidine or glutamic acid) yielded EPO muteins with biological activities similar to wild type EPO, whereas other amino acid changes (to alanine or glycine) resulted in EPO molecules with severely reduced activities (Delorme et al., 1992). The effect of changing serine-126 to cysteine was not studied. The in vivo bioactivity of S126V EPO was found to be similar to wild type EPO (Delorme et al., 1992).

The requirement for complex, N-linked carbohydrates containing terminal sialic acid residues for in vivo activity of EPO has limited commercial manufacture of the protein to mammalian cells. The important functions of the sialated N-linked sugars are to prevent protein aggregation, increase protein stability and prolong the circulating half-life of the protein. The terminal sialic acid residues prolong EPO's circulating half-life by masking the underlying galactose residues, which are recognized by specific receptors on hepatocytes and promote clearance of the asialated protein. EPO can be produced in insect cells and is N-glycosylated and fully active in vitro; its activity in vivo has not been reported (Wojchowski et al., 1987).

This example provides for the design of cysteine-added EPO variants and their use in preparing conjugates using cysteine-reactive PEGs and other cysteine-reactive moieties. Certain amino acids in EPO are non-essential for biological activity and can be mutated to

cysteine residues without altering the normal disulfide binding pattern and overall conformation of the molecule. These amino acids are located in the A-B loop (amino acids 23-58 of the mature protein sequence), the B-C loop (amino acids 77-89 of the mature protein sequence), the C-D loop (amino acids 108-131 of the mature protein sequence), proximal to helix A (amino acids 1-8) and distal to helix D (amino acids 153-166 of the mature protein sequence). Also contemplated as preferred sites for adding cysteine residues are at the N-terminus or C-terminus of the protein sequence. Preferred sites for cysteine substitutions are the O-linked glycosylation site (serine-126) and the amino acids comprising the three N-linked glycosylation sites (N24, I25, T26, N38, I39, T40, N83, S84, S85). Glycosylation sites are attractive sites for introducing cysteine substitutions and attaching PEG molecules to EPO because (1) these sites are surface exposed; (2) the natural protein can tolerate bulky sugar groups at these positions; (3) the glycosylation sites are located in the putative loop regions and away from the receptor binding site (Wen et al., 1994); and (4) mutagenesis studies indicate these sites (at least individually) are not essential for in vitro or in vivo activity (Yamaguchi et al., 1991; Delorme et al., 1992). As discussed above, the local conformation of the region encompassing the O-glycosylation site region seems to be important for biological activity. Whether a cysteine substitution at position 126 affects biological activity has not been studied. The cysteine-29 to cysteine-33 disulfide bond is not necessary for biological activity of EPO because changing both residues to tyrosine simultaneously yielded a biologically active EPO protein (Boissel et al., 1993; Wen et al., 1994). A "free" cysteine can be created by changing either cysteine-29 or cysteine-33 to another amino acid. Preferred amino acid changes would be to serine or alanine. The remaining "free" cysteine (cysteine-29 or cysteine-33) would be a preferred site for covalently modifying the protein with cysteine-reactive moieties.

Bill et al. (1995) individually substituted cysteine for N24, N38 and N83 and reported that the muteins had greatly reduced in vitro biological activities (less than 20% of wild type activity). Bill et al. (1995) expressed the EPO variants as fusion proteins (fused to glutathionine-S-transferase) in bacteria. One aspect of the present invention is to provide expression systems in which the N24C, N38C and N83C EPO variants will have in vitro biological activities more similar to wild type EPO.

US patent 4,703,008 contemplates naturally occurring variants of EPO as well as amino acid substitutions that are present in EPO proteins of mammals. Ovine EPO contains a

cysteine residue at position 88 of the polypeptide chain. The inventor is unaware of any other naturally occurring human or animal cysteine variants of EPO in which the cysteine residue occurs in the polypeptide regions disclosed herein as being useful for generating cysteine-added EPO variants. Patent 4,703,008 specifically teaches away from cysteine-added EPO variants by suggesting that expression of EPO might be improved by deleting cysteine residues or substituting naturally occurring cysteine residues with serine or histidine residues.

The mature protein form of EPO can contain 165 or 166 amino acids because of post-translational removal of the C-terminal arginine. Asp-165 is the C-terminus of the 165 amino acid form and Agr-166 is the C-terminal amino acid of the 166 amino acid form. The cysteine substitution and insertion mutations described herein can comprise either the 165 or 166 amino acid forms of mature EPO.

A cDNA encoding EPO can be cloned using the polymerase chain reaction (PCR) technique from the human HepG2 or Hep3B cell lines, which are known to express EPO when treated with hypoxia or cobalt chloride (Wen et al., 1993) and are available from the American Type Culture Collection (ATCC). Cysteine mutations can be introduced into the cDNA by standard phage, plasmid or PCR mutagenesis procedures as described for GH. As described above, the preferred sites for introduction of cysteine substitution mutations are in the A-B loop, the B-C loop, the C-D loop and the region proximal to helix A and distal to helix D. The most preferred sites in these regions are the N- and O-linked glycosylation sites: S126C; N24C; I25C, T26C; N38C; I39C, T40C; N83C, S 84C and S85C. Other preferred sites for cysteine substitution mutagenesis are in the A-B loop, the B-C loop and the C-D loop, amino acids surrounding the glycosylation sites and the region of the protein proximal to helix A and distal to helix D (Boissel et al., 1993; Wen et al., 1994). Other preferred sites for cysteine substitutions in these regions are: A1, P2, P3, R4, D8, S9, T27, G28, A30, E31, H32, S34, N36, D43, T44, K45, N47, A50, K52, E55, G57, Q58, G77, Q78, A79, Q86, W88, E89, T107, R110, A111, G113, A114, Q115, K116, E117, A118, S120, P121, P122, D123, A124, A125, A127, A128, T132, K154, T157, G158, E159, A160, T163, G164, D165 and R166. Cysteine residues also can be introduced proximal to the first amino acid of the mature protein, i.e., proximal to A1, or distal to the final amino acid in the mature protein, i.e., distal to D165 or R166. Other variants in which cys-29 or cys-33 have been replaced with other amino acids, preferably serine or alanine, also are provided.

Wild type EPO and EPO muteins can be expressed using insect cells to determine whether the cysteine-added muteins are biologically active. DNAs encoding EPO/EPO muteins can be cloned into the Baculovirus expression vector pVL1392 (available from Invitrogen, Inc. and Sigma Corporation (St. Louis, MO) and used to infect insect cells.

5 Recombinant Baculoviruses producing EPO can be identified by Western blots of infected insect cell conditioned media using polyclonal anti-human EPO antiserum (available from R&D Systems). The secreted EPO mutein proteins can be purified by conventional chromatographic procedures well known to those of skill in the art. Protein concentrations can be determined using commercially available protein assay kits or ELISA assay kits
10 (available from R&D Systems and Bio-Rad Laboratories).

Purified EPO and EPO muteins can be tested in cell proliferation assays using EPO-responsive cell lines such as UT7-epo (Wen et al., 1994) or TF1 (available from the ATCC) to measure specific activities of the proteins. Cells can be plated in 96-well microtiter plates with various concentrations of EPO. Assays should be performed in triplicate. After 1-3
15 days in culture, cell proliferation can be measured by ³H-thymidine incorporation, as described above for GH. The concentration of protein giving half-maximal stimulation (EC₅₀) can be determined for each mutein. Assays should be performed at least three times for each mutein, with triplicate wells for each data point. EC₅₀ values can be used to compare the relative potencies of the muteins. Alternatively, cell proliferation in response to
20 added EPO muteins can be analyzed using an MTT dye-exclusion assay (Komatsu et al. 1991). Proteins displaying similar optimal levels of stimulation and EC₅₀ values comparable to or greater than wild type EPO are preferable.

The above studies confirm identification of amino acid residues in EPO that can be changed to cysteine residues and retain biological activity. Muteins that retain activity can be
25 PEGylated using a cysteine-reactive 8 kDa PEG-maleimide as described above for GH muteins. Wild type EPO should be used as a control since it should not react with the cysteine-reactive PEG under identical partial reduction conditions. The lowest amount of PEG that gives significant quantities of mono-PEGylated product without giving di-PEGylated product should be considered optimum. Mono-PEGylated protein can be purified
30 from non-PEGylated protein and unreacted PEG by size-exclusion or ion exchange chromatography. The purified PEGylated proteins should be tested in the cell proliferation assay described above to determine their bioactivities.

One or more of the PEGylated EPO muteins that retain in vitro bioactivity, are candidates for testing in animal disease models. PEGylation of the protein at the proper amino acid can be determined as described for GH.

In vivo testing of PEGylated EPO muteins expressed using insect cells may require
5 that they be re-engineered for expression in mammalian cells to ensure proper glycosylation. PEG-EPO candidates produced using insect cells can be tested in the animal models described below to determine if they are active in vivo and whether they are as active as PEG-EPO produced using mammalian cell expression systems. For expression in mammalian cells, the EPO muteins can be subcloned into commercially available eukaryotic expression vectors
10 and used to stably transform Chinese Hamster Ovary (CHO) cells (available from the ATCC). Sublines can be screened for EPO expression using ELISA assays. Sufficient quantities of the insect cell- and mammalian cell-produced EPO muteins can be prepared to compare their biological activities in animal anemia models.

In vivo bioactivities of the EPO muteins can be tested using the artificial
15 polycythemia or starved rodent models (Cotes and Bangham., 1961; Goldwasser and Gross., 1975). In the starved rodent model, rats are deprived of food on day one and treated with test samples on days two and three. On day four, rats receive an injection of radioactive iron-59. Approximately 18h later, rats are anesthetized and blood samples drawn. The percent conversion of labeled iron into red blood cells is then determined. In the artificial
20 polycythemia model, mice are maintained in a closed tank and exposed for several days to hypobaric air. The animals are then brought to normal air pressure. Red blood cell formation is suppressed for several days. On day four or six after return to normal air pressure, mice are injected with erythropoietin or saline. Mice receive one injection per day for one to two days. One day later the animals receive an intravenous injection of labeled iron-59. The mice are
25 euthanized 20 h later and the amount of labeled iron incorporated into red blood cells determined. EPO stimulates red blood cell formation in both models as measured by a dose-dependent increase in labeled iron incorporated into red blood cells. In both models different dosing regimens and different times of injections can be studied to determine if PEG-EPO is biologically active and/or more potent and produces longer acting effects than natural EPO.

30

Example 3

Alpha Interferon

Alpha interferon is produced by leukocytes and has antiviral, anti-tumor and immunomodulatory effects. There are at least 20 distinct alpha interferon genes that encode proteins that share 70% or greater amino acid identity. Amino acid sequences of the known alpha interferon species are given in Blatt et al. (1996). A "consensus" interferon that incorporates the most common amino acids into a single polypeptide chain has been described (Blatt et al., 1996). A hybrid alpha interferon protein may be produced by splicing different parts of alpha interferon proteins into a single protein (Horisberger and Di Marco, 1995). Some alpha interferons contain N-linked glycosylation sites in the region proximal to helix A and near the B-C loop (Blatt et al. 1966). The alpha 2 interferon protein (SEQ ID NO: 3) contains four cysteine residues that form two disulfide bonds. The cys1-cys98 disulfide bond (cys1-cys99 in some alpha interferon species such as alpha 1; SEQ ID NO: 4) is not essential for activity. The alpha 2-interferon protein does not contain any N-linked glycosylation sites. The crystal structure of alpha interferon has been determined (Radhakrishnan et al., 1996).

This example provides cysteine added variants in the region proximal to the A helix, distal to the E helix, in the A-B loop, in the B-C loop, in the C-D loop and in the D-E loop. Preferred sites for the introduction of cysteine residues in these regions of the alpha interferon-2 species are: D2, L3, P4, Q5, T6, S8, Q20, R22, K23, S25, F27, S28, K31, D32, R33, D35, G37, F38, Q40, E41, E42, F43, G44, N45, Q46, F47, Q48, K49, A50, N65, S68, T69, K70, D71, S72, S73, A74, A75, D77, E78, T79, Y89, Q90, Q91, N93, D94, E96, A97, Q101, G102, G104, T106, E107, T108, P109, K112, E113, D114, S115, K131, E132, K133, K134, Y135, S136, A139, S152, S154, T155, N156, L157, Q158, E159, S160, L161, R162, S163, K164, E165. Variants in which cysteine residues are introduced proximal to the first amino acid of the mature protein, i.e., proximal to C1, or distal to the final amino acid in the mature protein, i.e., distal to E165 are provided. Other variants in which cys-1 or cys-98 (cys-99 in some alpha interferon species) have been replaced with other amino acids, preferably serine or alanine, also are provided. Other variants in which Cys-1 has been deleted (des Cys-1) also are provided. The cysteine variants may be in the context of any naturally occurring or non-natural alpha interferon sequence, e.g., consensus interferon or interferon protein hybrids. Some naturally occurring alpha interferon species (e.g., alpha interferon-1) contain a naturally occurring "free" cysteine. In such interferon species the naturally occurring free cysteine can be changed to another amino acid, preferably serine or alanine.

This example also provides cysteine variants of other alpha interferon species, including consensus interferon, at equivalent sites in these proteins. The alignment of the alpha interferon-2 species with other known alpha interferon species and consensus interferon is given in Blatt et al. (1996). The crystal structure of alpha interferon -2 has been determined
5 by Rhadhakrishnan et al. (1996). Lydon et al (1985) found that deletion of the first four amino acids from the N-terminus of alpha interferon did not affect biological activity. Valenzuela et al (1985) found that substitution of Phe-47 in alpha interferon-2 with Cys, Tyr or Ser did not alter biological activity of the protein. Cys-1 and Cys-98 have been changed individually to glycine and serine, respectively, without altering biological activity of the
10 protein (DeChiara et al, 1986).

DNA sequences encoding alpha interferon-2 can be amplified from human genomic DNA, since alpha interferon genes do not contain introns (Pestka et al., 1987). The DNA sequence of alpha interferon-2 is given in Goeddel et al. (1980). Alternatively, a cDNA for alpha interferon-2 can be isolated from human lymphoblastoid cell lines that are known to
15 express alpha interferon spontaneously or after exposure to viruses (Goeddel et al., 1980; Pickering et al., 1980). Many of these cell lines are available from the American Type Culture Collection (Rockville, MD). Specific mutations can be introduced into the alpha interferon sequence using plasmid-based site-directed mutagenesis kits (e.g., Quick-Change Mutagenesis Kit, Stratagene, Inc.), phage mutagenesis strategies or employing PCR
20 mutagenesis as described for GH.

Alpha interferon has been successfully produced in E. coli as an intracellular protein (Tarnowski et al., 1986; Thatcher and Panayotatos, 1986). Similar procedures can be used to express alpha interferon muteins. Plasmids encoding alpha interferon or alpha interferon muteins can be cloned into an E. coli expression vector such as pET15b (Novagene, Inc.) that
25 uses the strong T7 promoter or pCYB1 (New England BioLabs, Beverly, MA) that uses the TAC promoter. Expression of the protein can be induced by adding IPTG to the growth media.

Recombinant alpha interferon expressed in E. coli is sometimes soluble and sometimes insoluble (Tarnowski et al., 1986; Thatcher and Panayotatos, 1986). Insolubility
30 appears to be related to the degree of overexpression of the protein. Insoluble alpha interferon proteins can be recovered as inclusion bodies and renatured to a fully active conformation following standard oxidative refolding protocols (Thatcher and Panayotatos, 1986; Cox et al.,

1994). The alpha interferon proteins can be purified further using other chromatographic methods such as ion-exchange, hydrophobic interaction, size-exclusion and reversed phase resins (Thatcher and Panayotaos, 1986). Protein concentrations can be determined using commercially available protein assay kits (Bio-Rad Laboratories).

5 If E. coli expression of alpha interferon muteins is not successful, one can express the proteins in insect cells as secreted proteins as described for GH. The proteins can be modified to contain the natural alpha interferon signal sequence (Goeddell et al., 1980) or the honeybee mellitin signal sequence (Invitrogen, Inc.) to promote secretion of the proteins. Alpha interferon and alpha interferon muteins can be purified from conditioned media using
10 conventional chromatography procedures. Antibodies to alpha interferon can be used in conjunction with Western blots to localize fractions containing the alpha interferon proteins during chromatography. Alternatively, fractions containing alpha interferon proteins can be identified using ELISAs.

 Bioactivities of alpha interferon and alpha interferon muteins can be measured using
15 an in vitro viral plaque reduction assay (Ozes et al., 1992; Lewis, 1995). Human HeLa cells can be plated in 96-well plates and grown to near confluency at 37°C. The cells are then washed and treated for 24 hour with different concentrations of each alpha interferon preparation. Controls should include no alpha interferon and wild type alpha interferon (commercially available from Endogen, Inc., Woburn, MA). A virus such as Vesicular
20 stomatitis virus (VSV) or encephalomyocarditis virus (EMCV) is added to the plates and the plates incubated for a further 24-48 hours at 37°C. Additional controls should include samples without virus. When 90% or more of the cells have been killed in the virus-treated, no alpha interferon control wells (determined by visual inspection of the wells), the cell monolayer are stained with crystal violet and absorbance of the wells read using a microplate
25 reader. Alternatively, the cell monolayers can be stained with the dye MTT (Lewis, 1995). Samples should be analyzed in duplicate or triplicate. EC₅₀ values (the amount of protein required to inhibit the cytopathic effect of the virus by 50%) can be used to compare the relative potencies of the proteins. Wild type alpha interferon-2 protects cells from the cytopathic effects of VSV and EMCV and has a specific activity of approximately 2x10⁸
30 units/mg in this assay (Ozes et al., 1992). Alpha interferon muteins displaying EC₅₀ values comparable to wild type Alpha Interferon are preferable.

Alpha interferon muteins that retain activity can be PEGylated using procedures similar to those described for GH. Wild type alpha interferon-2 can serve as a control since it should not PEGylate under similar conditions. The lowest amount of PEG that gives significant quantities of mono-pegylated product without giving di-pegylated product should be considered optimum. Mono-PEGylated protein can be purified from non-PEGylated protein and unreacted PEG by size-exclusion or ion exchange chromatography. The purified PEGylated proteins can be tested in the viral plaque reduction bioassay described above to determine their bioactivities. PEGylated alpha interferon proteins with bioactivities comparable to wild type alpha interferon are preferable. Mapping the PEG attachment site and determination of pharmacokinetic data for the PEGylated protein can be performed as described for GH.

In vivo bioactivities of the PEG-alpha interferon muteins can be tested using tumor xenograft models in nude mice and viral infection models (Balkwill, 1986; Fish et al., 1986). Since PEG-alpha interferon bioactivity may be species-specific, one should confirm activity of the PEGylated protein using appropriate animal cell lines in in vitro virus plaque reduction assays, similar to those described above. Next, one should explore the effects of different dosing regimens and different times of injections to determine if PEG-alpha interferon is more potent and produces longer lasting effects than non-PEGylated alpha interferon.

The novel alpha interferon-derived molecules of this example can be formulated and tested for activity essentially as set forth in Examples 1 and 2, substituting, however, the appropriate assays and other considerations known in the art related to the specific proteins of this example.

Example 4

Beta Interferon

Beta interferon is produced by fibroblasts and exhibits antiviral, antitumor and immunomodulatory effects. The single-copy beta interferon gene encodes a preprotein that is cleaved to yield a mature protein of 166 amino acids (Taniguchi et al. 1980; SEQ ID NO: 5). The protein contains three cysteines, one of which, cysteine-17, is "free", i.e., it does not participate in a disulfide bond. The protein contains one N-linked glycosylation site. The crystal structure of the protein has been determined (Karpusas et al. 1997)

This example provides cysteine-added variants at any of the three amino acids that comprise the N-linked glycosylation sites, i.e., N80C, E81C or T82C. This example also provides cysteine-added variants in the region proximal to the A helix, distal to the E helix, in the A-B loop, in the B-C loop, in the C-D loop and in the D-E loop. Preferred sites for introduction of cysteine residues in these regions are: M1, S2, Y3, N4, L5, Q23, N25, G26, R27, E29, Y30, K33, D34, R35, N37, D39, E42, E43, K45, Q46, L47, Q48, Q49, Q51, K52, E53, A68, F70, R71, Q72, D73, S74, S75, S76, T77, G78, E107, K108, E109, D110, F111, T112, R113, G114, K115, L116, A135, K136, E137, K138, S139, I157, N158, R159, L160, T161, G162, Y163, L164, R165 and N166. Variants in which cysteine residues are introduced proximal to the first amino acid of the mature protein, i.e., proximal to M1, or distal to the final amino acid in the mature protein, i.e., distal to N166 also are provided.

These variants are produced in the context of the natural protein sequence or a variant protein in which the naturally occurring "free" cysteine residue (cysteine-17) has been changed to another amino acid, preferably serine or alanine.

The novel beta interferon-derived molecules of this example can be formulated and tested for activity essentially as set forth in Examples 1 and 2, substituting, however, the appropriate assays and other considerations known in the art related to the specific proteins of this example.

Example 5

Granulocyte Colony-Stimulating Factor (G-CSF)

G-CSF is a pleuripotent cytokine that stimulates the proliferation, differentiation and function of granulocytes. The protein is produced by activated monocytes and macrophages. The amino acid sequence of G-CSF (SEQ ID NO: 6) is given in Souza et al. (1986), Nagata et al. (1986a, b) and US patent 4,810,643 all incorporated herein by reference. The human protein is synthesized as a preprotein of 204 or 207 amino acids that is cleaved to yield mature proteins of 174 or 177 amino acids. The larger form has lower specific activity than the smaller form. The protein contains five cysteines, four of which are involved in disulfide bonds. Cysteine-17 is not involved in a disulfide bond. Substitution of cysteine-17 with serine yields a mutant G-CSF protein that is fully active (US patent 4,810,643). The protein is O-glycosylated at threonine-133 of the mature protein.

This example provides a cysteine-added variant at threonine-133. This example provides other cysteine-added variants in the region proximal to helix A, distal to helix D, in the A-B loop, B-C loop and C-D loop. Preferred sites for introduction of cysteine substitutions in these regions are: T1, P2, L3, G4, P5, A6, S7, S8, L9, P10, Q11, S12, T38, K40, S53, G55, W58, A59, P60, S62, S63, P65, S66, Q67, A68, Q70, A72, Q90, A91, E93, G94, S96, E98, G100, G125, M126, A127, A129, Q131, T133, Q134, G135, A136, A139, A141, S142, A143, Q145, Q173 and P174. Variants in which cysteine residues are introduced proximal to the first amino acid of the mature protein, i.e., proximal to T1, or distal to the final amino acid in the mature protein, i.e., distal to P174 are provided. These variants are provided in the context of the natural protein sequence or a variant protein in which the naturally occurring "free" cysteine residue (cysteine-17) has been changed to another amino acid, preferably serine or alanine.

A cDNA encoding human G-CSF can be purchased from R&D Systems (Minneapolis, MN) or amplified using PCR from mRNA isolated from human carcinoma cell lines such as 5637 and U87-MG known to express G-CSF constitutively (Park et al., 1989; Nagata, 1994). These cell lines are available from the American Type Culture collection (Rockville, MD). Specific mutations can be introduced into the G-CSF sequence using plasmid-based site-directed mutagenesis kits (e.g., Quick-Change Mutagenesis Kit, Stratagene, Inc.), phage mutagenesis methods or employing PCR mutagenesis as described for GH.

G-CSF has been successfully produced in *E. coli* as an intracellular protein (Souza et al., 1986). One can employ similar procedures to express G-CSF and G-CSF muteins. Plasmids encoding G-CSF or G-CSF muteins can be cloned into an *E. coli* expression vector such as pET15b (available from Novagen, Inc., Madison, WI) that uses the strong T7 promoter or pCYB1 (available from New England BioLabs, Beverly, MA) that uses the strong TAC promoter. Expression of the protein can be induced by adding IPTG to the growth media. Recombinant G-CSF expressed in *E. coli* is insoluble and can be recovered as inclusion bodies. The protein can be renatured to a fully active conformation following standard oxidative refolding protocols (Souza et al., 1986; Lu et al., 1992; Cox et al., 1994). Similar procedures can be used to refold cysteine muteins. The proteins can be purified further using other chromatographic methods such as ion exchange, hydrophobic interaction, size-exclusion and reversed phase resins (Souza et al., 1986; Kuga et al., 1989; Lu et al.,

1992). Protein concentrations can be determined using commercially available protein assay kits (Bio-Rad Laboratories).

If E. coli expression of G-CSF or G-CSF muteins is not successful, one can express G-CSF and G-CSF muteins in insect cells as secreted proteins as described for GH. The
5 proteins can be modified to contain the natural G-CSF signal sequence (Souza et al., 1986; Nagata et al., 1986a; Nagata et al., 1986b) or the honeybee mellitin signal sequence (Invitrogen, Inc., Carlsbad, CA) to promote secretion of the proteins. G-CSF and G-CSF muteins can be purified from conditioned media using conventional chromatography procedures. Antibodies to G-CSF can be used in conjunction with Western blots to localize
10 fractions containing the G-CSF proteins during chromatography. Alternatively, fractions containing G-CSF proteins can be identified using ELISAs.

G-CSF muteins also can be expressed in mammalian cells as described for erythropoietin in Example 2.

Bioactivities of G-CSF and the G-CSF muteins can be measured using an in vitro cell
15 proliferation assay. The mouse NFS-60 cell line and the human AML-193 cell line can be used to measure G-CSF bioactivity (Tsuchiya et al., 1986; Lange et al., 1987; Shirafuji et al., 1989). Both cell lines proliferate in response to human G-CSF. The AML-193 cell line is preferable since it is of human origin, which eliminates the possibility of a false conclusion resulting from species differences. The NFS60 cell lines proliferates in response to G-CSF
20 with a half-maximal effective concentration (EC_{50}) of 10-20 picomolar. Purified G-CSF and G-CSF muteins can be tested in cell proliferation assays using these cell lines to determine specific activities of the proteins, using published methods (Tsuchiya et al., 1986; Lange et al., 1987; Shirafuji et al., 1989). Cells can be plated in 96-well tissue culture dishes with different concentrations of G-CSF or G-CSF muteins. After 1-3 days at 37°C in a humidified
25 tissue culture incubator, proliferation can be measured by 3H -thymidine incorporation as described for GH. Assays should be performed at least three times for each mutein using triplicate wells for each data point. EC_{50} values can be used to compare the relative potencies of the muteins. G-CSF muteins displaying similar optimal levels of stimulation and EC_{50} values comparable to wild type G-CSF are preferable.

30 G-CSF muteins that retain activity can be PEGylated using procedures similar to those described for GH. Wild type G-CSF and ser-17 G-CSF can serve as controls since they should not PEGylate under similar conditions. The lowest amount of PEG that gives

significant quantities of mono-PEGylated product without giving di-PEGylated product should be considered optimum. Mono-PEGylated protein can be purified from non-PEGylated protein and unreacted PEG by size-exclusion or ion exchange chromatography. The purified PEGylated proteins can be tested in the cell proliferation assay described above
5 to determine their bioactivities.

The PEG site in the protein can be mapped using procedures similar to those described for GH. Pharmacokinetic data for the PEGylated proteins can be obtained using procedures similar to those described for GH.

Initial studies to demonstrate in vivo efficacy of PEG-G-CSF can be done in normal
10 Sprague-Dawley rats, which can be purchased from Charles River. Groups of rats should receive single subcutaneous or intravenous injections of various doses of G-CSF, PEG-G-CSF or placebo. Animals should be sacrificed at daily intervals for up to a week for determination of neutrophil and total white blood cell counts in peripheral blood. Other blood cell types (platelets and red blood cells) can be measured to demonstrate cell specificity.

15 The efficacy of PEG-G-CSF can be tested in a rat neutropenia model. Neutropenia can be induced by treatment with cyclophosphamide, which is a commonly used chemotherapeutic agent that is myelosuppressive. G-CSF accelerates recovery of normal neutrophil levels in cyclophosphamide-treated animals (Kubota et al., 1990). Rats receive an injection of cyclophosphamide on day 0 to induce neutropenia. The animals are then be
20 divided into different groups, which will receive subcutaneous injections of G-CSF, PEG-G-CSF or placebo. Neutrophil and total white blood cell counts in peripheral blood should be measured daily until they return to normal levels. Initially one should confirm that G-CSF accelerates recovery from neutropenia when injected daily. Next, one should explore the effects of different dosing regimens and different times of injections to determine if PEG-G-
25 CSF is more potent and produces longer lasting effects than non-PEGylated G-CSF.

The novel GCSF-derived molecules of this example can be formulated and tested for activity essentially as set forth in Examples 1 and 2, substituting, however, the appropriate assays and other considerations known in the art related to the specific proteins of this example.

30

Example 6

Thrombopoietin (TPO)

Thrombopoietin stimulates the development of megakaryocyte precursors of platelets. The amino acid sequence of TPO (SEQ ID NO: 7) is given in Bartley et al.(1994), Foster et al.(1994), de Sauvage et al.(1994), each incorporated herein by reference.

The protein is synthesized as a 353 amino acid precursor protein that is cleaved to
5 yield a mature protein of 332 amino acids. The N-terminal 154 amino acids have homology with EPO and other members of the GH supergene family. The C-terminal 199 amino acids do not share homology with any other known proteins. The C-terminal region contains six N-linked glycosylation sites and multiple O-linked glycosylation sites (Hoffman et al., 1996). O-linked glycosylation sites also are found in the region proximal to the A helix, in the A-B
10 loop, and at the C-terminus of Helix C (Hoffman et al., 1996). A truncated TPO protein containing only residues 1-195 of the mature protein is fully active in vitro (Bartley et al., 1994).

This example provides cysteine-added variants at any of the amino acids that comprise the N-linked glycosylation sites and the O-linked glycosylation sites. This example also
15 provides cysteine-added variants in the region proximal to the A helix, distal to the D helix, in the A-B loop, in the B-C loop, and in the C-D loop.

Preferred sites for introduction of cysteine residues are: S1, P2, A3, P4, P5, A6, T37, A43, D45, S47, G49, E50, K52, T53, Q54, E56, E57, T58, A76, A77, R78, G79, Q80, G82, T84, S87, S88, G109, T110, Q111, P113, P114, Q115, G116, R117, T118, T119, A120,
20 H121, K122, G146, G147, S148, T149, A155, T158, T159, A160, S163, T165, S166, T170, N176, R177, T178, S179, G180, E183, T184, N185, F186, T187, A188, S189, A190, T192, T193, G194, S195, N213, Q214, T215, S216, S218, N234, G235, T236, S244, T247, S254, S255, T257, S258, T260, S262, S272, S274, T276, T280, T291, T294, S307, T310, T312, T314, S315, N319, T320, S321, T323, S325, Q326, N327, L328, S329, Q330, E331 and
25 G332. Variants in which cysteine residues are introduced proximal to the first amino acid of the mature protein, i.e., proximal to S1, or distal to the final amino acid in the mature protein, i.e., distal to G332 are provided. The cysteine-added variants are provided in the context of the natural human protein or a variant protein that is truncated between amino acids 147 and the C-terminus of the natural protein, G332. Variants in which cysteine residues are added
30 distal to the final amino acid of a TPO protein that is truncated between amino acids 147 and 332 also are provided.

The novel TPO-derived molecules of this example can be formulated and tested for activity essentially as set forth in Examples 1 and 2, substituting, however, the appropriate assays and other considerations known in the art related to the specific proteins of this example.

5

Example 7

Granulocyte-Macrophage Colony Stimulating Factor (GM-CSF)

GM-CSF stimulates the proliferation and differentiation of various hematopoietic cells, including neutrophil, monocyte, eosinophil, erythroid, and megakaryocyte cell lineages.

10 The amino acid sequence of human GM-CSF (SEQ ID NO: 8) is given in Cantrell et al. (1985) and Lee et al (1985) both incorporated herein by reference.

GM-CSF is produced as a 144 amino acid preprotein that is cleaved to yield a mature 127 amino acid protein. The mature protein has two sites for N-linked glycosylation. One site is located at the C-terminal end of Helix A; the second site is in the A-B loop.

15 This example provides cysteine-added variants at any of the amino acids that comprise the N-linked glycosylation sites, i.e., N27C, L28C, S29C, N37C, E38C and T39C. This example also provides cysteine-added variants in the region proximal to the A helix, distal to the D helix, in the A-B loop, in the B-C loop, and in the C-D loop. Preferred sites for introduction of cysteine substitutions in these regions are: A1, P2, A3, R4, S5, P6, S7, P8, S9, 20 T10, Q11, R30, D31, T32, A33, A34, E35, E41, S44, E45, D48, Q50, E51, T53, Q64, G65, R67, G68, S69, L70, T71, K72, K74, G75, T91, E93, T94, S95, A97, T98, T102, I117, D120, E123, V125, Q126 and E127. Variants in which cysteine residues are introduced proximal to the first amino acid of the mature protein, i.e., proximal to A1, or distal to the final amino acid in the mature protein, i.e., distal to E127 are provided.

25 The novel GM-CSF-derived molecules of this example can be formulated and tested for activity essentially as set forth in Examples 1 and 2, substituting, however, the appropriate assays and other considerations known in the art related to the specific proteins of this example.

Example 8**IL-2**

IL-2 is a T cell growth factor that is synthesized by activated T cells. The protein stimulates clonal expansion of activated T cells. Human IL-2 is synthesized as a 153 amino acid precursor that is cleaved to yield a 133 amino acid mature protein (Tadatsugu et al., 1983; Devos et al., 1983; SEQ ID NO: 9).

The amino acid sequence of IL-2 is set forth in (Tadatsugu et al. 1983; Devos et al. 1983). The mature protein contains three cysteine residues, two of which form a disulfide bond. Cysteine-125 of the mature protein is not involved in a disulfide bond. Replacement of cysteine-125 with serine yields an IL-2 mutein with full biological activity (Wang et al., 1984). The protein is O-glycosylated at threonine-3 of the mature protein chain.

This example provides cysteine-added variants in the last four positions of the D helix, in the region distal to the D helix, in the A-B loop, in the B-C loop, and in the C-D loop. These variants are provided in the context of the natural protein sequence or a variant protein in which the naturally occurring "free" cysteine residue (cysteine-125) has been changed to another amino acid, preferably serine or alanine. Variants in which cysteine residues are introduced proximal to the first amino acid, i.e., A1, or distal to the final amino acid, i.e., T133, of the mature protein, also are provided.

The novel IL-2-derived molecules of this example can be formulated and tested for activity essentially as set forth in Examples 1 and 2, substituting, however, the appropriate assays and other considerations known in the art related to the specific proteins of this example.

Example 9**IL-3**

IL-3 is produced by activated T cells and stimulates the proliferation and differentiation of pleuripotent hematopoietic stem cells. The amino acid sequence of human IL-3 (SEQ ID NO: 10) is given in Yang et al. (1986); Dorssers et al. (1987) and Otsuka et al. (1988) all incorporated herein by reference. The protein contains two cysteine residues and two N-linked glycosylation sites. Two alleles have been described, resulting in isoforms having serine or proline at amino acid position 8 of the mature protein.

This example provides cysteine-added variants at any of the amino acids that comprise the N-linked glycosylation sites. This example also provides cysteine-added variants in the region proximal to the A helix, distal to the D helix, in the A-B loop, in the B-C loop, and in the C-D loop. Variants in which cysteine residues are introduced proximal to the first amino acid or distal to the final amino acid in the mature protein, also are provided.

The novel IL-3-derived molecules of this example can be formulated and tested for activity essentially as set forth in Examples 1 and 2, substituting, however, the appropriate assays and other considerations known in the art related to the specific proteins of this example.

Example 10

IL-4

IL-4 is a pleiotropic cytokine that stimulates the proliferation and differentiation of monocytes and T and B cells. IL-4 has been implicated in the process that leads to B cells secreting IgE, which is believed to play a role in asthma and atopy. The bioactivity of IL-4 is species specific. IL-4 is synthesized as a 153 amino acid precursor protein that is cleaved to yield a mature protein of 129 amino acids. The amino acid sequence of human IL-4 (SEQ ID NO:11) is given in Yokota et al. (1986) which is incorporated herein by reference. The protein contains six cysteine residues and two N-linked glycosylation sites. The glycosylation sites are located in the A-B and C-D loops.

This example provides cysteine-added variants at any of the amino acids that comprise the N-linked glycosylation sites. This example also provides cysteine-added variants in the region proximal to the A helix, distal to the D helix, in the A-B loop, in the B-C loop, and in the C-D loop. Variants in which cysteine residues are introduced proximal to the first amino acid, H1, or distal to the final amino acid, S129, of the mature protein are provided.

The novel IL-4-derived molecules of this example can be formulated and tested for activity essentially as set forth in Examples 1 and 2, substituting, however, the appropriate assays and other considerations known in the art related to the specific proteins of this example.

Example 11

IL-5

IL-5 is a differentiation and activation factor for eosinophils. The amino acid sequence of human IL-5 (SEQ ID NO: 12) is given in Yokota et al. (1987) which is incorporated herein by reference. The mature protein contains 115 amino acids and exists in solution as a disulfide-linked homodimer. The protein contains both O-linked and N-linked glycosylation sites.

This example provides cysteine-added variants in the region proximal to the A helix, distal to the D helix, in the A-B loop, in the B-C loop, and in the C-D loop. Variants in which cysteine residues are added proximal to the first amino acid or distal to the final amino acid in the mature protein are provided.

The novel IL-5-derived molecules of this example can be formulated and tested for activity essentially as set forth in Examples 1 and 2, substituting, however, the appropriate assays and other considerations known in the art related to the specific proteins of this example.

Example 12

IL-6

IL-6 stimulates the proliferation and differentiation of many cell types. The amino acid sequence of human IL-6 (SEQ ID NO: 13) is given in Hirano et al. (1986) which is incorporated herein by reference. IL-6 is synthesized as a 212 amino acid preprotein that is cleaved to generate a 184 amino acid mature protein. The mature protein contains two sites for N-linked glycosylation and one site for O-glycosylation, at T137, T138, T142 or T143.

This example provides cysteine-added variants at any of the amino acids that comprise the N-linked glycosylation sites and at the O-linked glycosylation site. Also provided are cysteine-added variants in the region proximal to the A helix, distal to the D helix, in the A-B loop, in the B-C loop, and in the C-D loop. Variants in which cysteine residues are added proximal to the first amino acid or distal to the final amino acid of the mature protein also are provided.

The novel IL-6-derived molecules of this example can be formulated and tested for activity essentially as set forth in Examples 1 and 2, substituting, however, the appropriate assays and other considerations known in the art related to the specific proteins of this example.

Example 13

IL-7

IL-7 stimulates proliferation of immature B cells and acts on mature T cells. The amino acid sequence of human IL-7 (SEQ ID NO: 14) is given in Goodwin et al. (1989) which is incorporated herein by reference. The protein is synthesized as a 177 amino acid preprotein that is cleaved to yield a 152 amino acid mature protein that contains three sites for N-linked glycosylation.

This example provides cysteine-added variants at any of the amino acids that comprise the N-linked glycosylation sites. This example also provides cysteine-added variants in the region proximal to the A helix, distal to the D helix, in the A-B loop, in the B-C loop, and in the C-D loop.

The novel IL-7-derived molecules of this example can be formulated and tested for activity essentially as set forth in Examples 1 and 2, substituting, however, the appropriate assays and other considerations known in the art related to the specific proteins of this example. Variants in which cysteine residues are added proximal to the first amino acid or distal to the final amino acid of the mature protein also are provided.

Example 14

IL-9

IL-9 is a pleiotropic cytokine that acts on many cell types in the lymphoid, myeloid and mast cell lineages. IL-9 stimulates the proliferation of activated T cells and cytotoxic T lymphocytes, stimulates proliferation of mast cell precursors and synergizes with erythropoietin in stimulating immature red blood cell precursors. The amino acid sequence of human IL-9 (SEQ ID NO: 15) is given in Yang et al. (1989) which is incorporated herein by reference. IL-9 is synthesized as a precursor protein of 144 amino acids that is cleaved to yield a mature protein of 126 amino acids. The protein contains four potential N-linked glycosylation sites.

This example provides cysteine-added variants at any of the three amino acids that comprise the N-linked glycosylation sites. This example also provides cysteine-added variants in the region proximal to the A helix, distal to the D helix, in the A-B loop, in the B-C loop, and in the C-D loop. Variants in which cysteine residues are added proximal to the first amino acid or distal to the final amino acid of the mature protein also are provided.

The novel IL-9-derived molecules of this example can be formulated and tested for activity essentially as set forth in Examples 1 and 2, substituting, however, the appropriate assays and other considerations known in the art related to the specific proteins of this example.

5

Example 15

IL-10

The amino acid sequence of human IL-10 (SEQ ID NO: 16) is given in Vieira et al. (1991) which is incorporated herein by reference. IL-10 is synthesized as a 178 amino acid precursor protein that is cleaved to yield a mature protein of 160 amino acids. IL-10 can function to activate or suppress the immune system. . The protein shares structural homology with the interferons, i.e., it contains five amphipathic helices. The protein contains one N-linked glycosylation site.

This example provides cysteine-added variants at any of the three amino acids comprising the N-linked glycosylation site. This example also provides cysteine-added variants in the region proximal to the A helix, distal to the E helix, in the A-B loop, in the B-C loop, in the C-D loop and in the D-E loop. Variants in which cysteine residues are added proximal to the first amino acid or distal to the final amino acid of the mature protein also are provided.

The novel IL-10-derived molecules of this example can be formulated and tested for activity essentially as set forth in Examples 1 and 2, substituting, however, the appropriate assays and other considerations known in the art related to the specific proteins of this example.

25

Example 16

IL-11

IL-11 is a pleiotropic cytokine that stimulates hematopoiesis, lymphopoiesis and acute phase responses. IL-11 shares many biological effects with IL-6. The amino acid sequence of human IL-11 (SEQ ID NO: 17) is given in Kawashima et al. (1991) and Paul et al. (1990) both incorporated herein by reference. IL-11 is synthesized as a precursor protein of 199 amino acids that is cleaved to yield a mature protein of 178 amino acids. There are no N-linked glycosylation sites in the protein.

30

This example provides cysteine-added variants in the region proximal to the A helix, distal to the D helix, in the A-B loop, in the B-C loop, and in the C-D loop. Variants in which cysteine residues are added proximal to the first amino acid or distal to the final amino acid of the mature protein also are provided.

The novel IL-11-derived molecules of this example can be formulated and tested for activity essentially as set forth in Examples 1 and 2, substituting, however, the appropriate assays and other considerations known in the art related to the specific proteins of this example.

Example 17

IL-12 p35

IL-12 stimulates proliferation and differentiation of NK cells and cytotoxic T lymphocytes. IL-12 exists as a heterodimer of a p35 subunit and a p40 subunit. The p35 subunit is a member of the GH supergene family. The amino acid sequence of the p35 subunit (SEQ ID NO: 18) is given in Gubler et al. (1991) and Wolf et al. (1991) both incorporated herein by reference. p35 is synthesized as a precursor protein of 197 amino acids and is cleaved to yield a mature protein of 175 amino acids. The protein contains 7 cysteine residues and three potential N-linked glycosylation sites.

This example provides cysteine-added variants at any of the three amino acids that comprise the three N-linked glycosylation sites. This example also provides cysteine-added variants in the region proximal to the A helix, distal to the D helix, in the A-B loop, in the B-C loop, and in the C-D loop. These variants are provided in the context of the natural protein sequence or a variant protein in which the naturally occurring “free” cysteine residue has been changed to another amino acid, preferably serine or alanine. Variants in which cysteine residues are added proximal to the first amino acid or distal to the final amino acid of the mature protein also are provided.

The novel IL-12 p35-derived molecules of this example can be formulated and tested for activity essentially as set forth in Examples 1 and 2, substituting, however, the appropriate assays and other considerations known in the art related to the specific proteins of this example.

Example 18

IL-13

IL-13 shares many biological properties with IL-4. The amino acid sequence of human IL-13 (SEQ ID NO: 19) is given in McKenzie et al. (1993) and Minty et al. (1993) both incorporated herein by reference. The protein is synthesized as a 132 amino acid precursor protein that is cleaved to yield a mature protein of 112 amino acids. The mature protein contains 5 cysteine residues and multiple N-linked glycosylation sites. A variant in which glutamine at position 78 is deleted due to alternative mRNA splicing has been described (McKenzie et al 1993)

This example provides cysteine-added variants at any of the three amino acids comprising the N-linked glycosylation sites. This example also provides cysteine-added variants in the region proximal to the A helix, distal to the D helix, in the A-B loop, in the B-C loop, and in the C-D loop. These variants are provided in the context of the natural protein sequence or a variant sequence in which the pre-existing "free" cysteine has been changed to another amino acid, preferably to alanine or serine. Variants in which cysteine residues are added proximal to the first amino acid or distal to the final amino acid of the mature protein also are provided.

The novel IL-13-derived molecules of this example can be formulated and tested for activity essentially as set forth in Examples 1 and 2, substituting, however, the appropriate assays and other considerations known in the art related to the specific proteins of this example.

Example 19

IL-15

IL-15 stimulates the proliferation and differentiation of T cells, NK cells, LAK cells and Tumor Infiltrating Lymphocytes. IL-15 may be useful for treating cancer and viral infections. The sequence of IL-15 (SEQ ID NO: 20) is given in Anderson et al. (1995) which is incorporated herein by reference. IL-15 contains two N-linked glycosylation sites, which are located in the C-D loop and C-terminal end of the D helix. IL-15 encodes a 162 amino acid preprotein that is cleaved to generate a mature 114 amino acid protein.

This example provides cysteine-added variants at any of the three amino acids comprising the N-linked glycosylation sites in the C-D loop or C-terminal end of the D helix.

This example also provides cysteine-added variants proximal to helix A, in the A-B loop, the B-C loop, the C-D loop or distal to helix D. Variants in which cysteine residues are added proximal to the first amino acid or distal to the final amino acid of the mature protein also are provided.

5 The novel IL-15-derived molecules of this example can be formulated and tested for activity essentially as set forth in Examples 1 and 2, substituting, however, the appropriate assays and other considerations known in the art related to the specific proteins of this example.

10

Example 20

Macrophage Colony Stimulating Factor (M-CSF):

M-CSF regulates the growth, differentiation and function of monocytes. The protein is a disulfide-linked homodimer. Multiple molecular weight species of M-CSF, which arise from differential mRNA splicing, have been described. The amino acid sequence of human
15 M-CSF and its various processed forms are given in Kawasaki et al (1985), Wong et al. (1987) and Cerretti et al (1988) which are incorporated herein by reference. Cysteine-added variants can be produced following the general teachings of this application and in accordance with the examples set forth herein.

20 The novel MCSF-derived molecules of this example can be formulated and tested for activity essentially as set forth in Examples 1 and 2, substituting, however, the appropriate assays and other considerations known in the art related to the specific proteins of this example.

25

Example 21

Oncostatin M:

Oncostatin M is a multifunctional cytokine that affects the growth and differentiation of many cell types. The amino acid sequence of human oncostatin M (SEQ ID NO: 21) is given in Malik et al. (1989) which is incorporated herein by reference. Oncostatin M is produced by activated monocytes and T lymphocytes. Oncostatin M is synthesized as a 252
30 amino acid preprotein that is cleaved sequentially to yield a 227 amino acid protein and then a 196 amino acid protein (Linsley et al., 1990). The mature protein contains O-linked

glycosylation sites and two N-linked glycosylation sites. The protein is O-glycosylated at T160, T162 and S165. The mature protein contains five cysteine residues.

This example provides cysteine-added variants at either of the three amino acids comprising the N-linked glycosylation sites or at the amino acids that comprise the O-linked glycosylation sites. This example also provides cysteine-added variants proximal to helix A, in the A-B loop, in the B-C loop, in the C-D loop or distal to helix D. These variants are provided in the context of the natural protein sequence or a variant sequence in which the pre-existing "free" cysteine has been changed to another amino acid, preferably to alanine or serine. Variants in which cysteine residues are added proximal to the first amino acid or distal to the final amino acid of the mature protein also are provided.

The novel Oncostatin M-derived molecules of this example can be formulated and tested for activity essentially as set forth in Examples 1 and 2, substituting, however, the appropriate assays and other considerations known in the art related to the specific proteins of this example.

Example 22

Ciliary Neurotrophic Factor (CNTF):

The amino acid sequence of human CNTF (SEQ ID NO: 22) is given in Lam et al. (1991) which is incorporated herein by reference. CNTF is a 200 amino acid protein that contains no glycosylation sites or signal sequence for secretion. The protein contains one cysteine residue. CNTF functions as a survival factor for nerve cells.

This example provides cysteine-added variants proximal to helix A, in the A-B loop, the B-C loop, the C-D loop or distal to helix D. These variants are provided in the context of the natural protein sequence or a variant sequence in which the pre-existing "free" cysteine has been changed to another amino acid, preferably to alanine or serine. Variants in which cysteine residues are added proximal to the first amino acid or distal to the final amino acid of the mature protein also are provided.

The novel CNTF-derived molecules of this example can be formulated and tested for activity essentially as set forth in Examples 1 and 2, substituting, however, the appropriate assays and other considerations known in the art related to the specific proteins of this example.

Example 23

Leukemia Inhibitory Factor (LIF)

The amino acid sequence of LIF (SEQ ID NO: 23) is given in Moreau et al. (1988) and Gough et al. (1988) both incorporated herein by reference. The human gene encodes a
5 202 amino acid precursor that is cleaved to yield a mature protein of 180 amino acids. The protein contains six cysteine residues, all of which participate in disulfide bonds. The protein contains multiple O- and N-linked glycosylation sites. The crystal structure of the protein was determined by Robinson et al. (1994). The protein affects the growth and differentiation of many cell types.

10 This example provides cysteine-added variants at any of the three amino acids comprising the N-linked glycosylation sites or the O-linked glycosylation site. Also provided are cysteine-added variants proximal to helix A, in the A-B loop, the B-C loop, the C-D loop or distal to helix D. Variants in which cysteine residues are added proximal to the first amino acid or distal to the final amino acid of the mature protein also are provided.

15 The novel LIF-derived molecules of this example can be formulated and tested for activity essentially as set forth in Examples 1 and 2, substituting, however, the appropriate assays and other considerations known in the art related to the specific proteins of this example.

20 All of the documents cited herein are incorporated herein by reference.

The protein analogues disclosed herein can be used for the known therapeutic uses of the native proteins in essentially the same forms and doses all well known in the art.

While the exemplary preferred embodiments of the present invention are described herein with particularity, those having ordinary skill in the art will recognize changes,
25 modifications, additions, and applications other than those specifically described herein, and may adapt the preferred embodiments and methods without departing from the spirit of this invention.

References:

30 Abdel-Meguid, S.S., Shieh, h.-S., Smith W.W., Dayringer, H.E., Violand, B.N. and Bentle, L.A. (1987) Proc. Natl. Acad. Sci. USA 84: 6434-6437.

- Abuchowski, A., Kazo, G.M., Verhoest, C.R., van Es, T., Kafkewitz, D., Nucci, M.L., Viau, A.T. and Davis, F.F. (1984) *Cancer Biochem. Biophys.* 7: 175-186.
- Aggarwal, B.B. (1998) *Human Cytokines. Handbook for Basic and Clinical Research. Volume III.* Blackwell Science, Malden, MA.
- 5 Aggarwal, B.B. and Gutterman, J.U. (1992) *Human Cytokines. Handbook for Basic and Clinical Research. Volume I.* Blackwell Scientific Publications, Cambridge, MA.
- Aggarwal, B.B. and Gutterman, J.U. (1996) *Human Cytokines. Handbook for Basic and Clinical Research. Volume II.* Blackwell Science, Cambridge, MA.
- Anderson, D.M., Johnson, L., Glaccum, M.B., Copeland, N.G., Gilbert, D.J., Jenkins, N.A.,
 10 Valentine, V., Kirstein, M.N., Shapiro, D.N., Morris, S.W., Grabstein, K. and cosman, D. (1995) *Genomics* 25: 701-706.
- Balkwill, F.R. (1986) *Methods Enzymology* 119: 649-657.
- Bartley, T.D., Bogenberger, J. et al., (1994) *Cell* 77: 1117-1124.
- Bazan, F. (1991) *Immunology Today* 11: 350-354.
- 15 Bazan, J.F. (1992) *Science* 257: 410-411.
- Becker, G.W. and Hsiung, H. M. (1986) *FEBS Letters* 204: 145-150.
- Bill, R.M., Winter, P.C., McHale, C.M., Hodges, V.M., Elder, G.E., Caley, J., Flitsch, S.L., Bicknell, R. Lappin, T.R.J. (1995) *Biochem. Biophys. Acta* 1261: 35-43.
- Bittorf, T., Jaster, R., Brock, J. (1993) *FEBS Letters* 336: 133-136.
- 20 Blatt, L.M., Davis, J.M., Klein, S.B. and Taylor, M.W. (1996) *Journal of Interferon and cytokine Research* 16: 489-499.
- Boissel, J.-P., Lee, W.-R., Presnell, S.R., Cohen, F.E. and Bunn, H.F. (1993) *J. Biol. Chem.* 268: 15983-15993.
- Butler, C.A. (1996) Lehman Brothers Technical Report "A Paradigm for Success".
- 25 Cantrell, M.A., Anderson, D., Cerretti, D.P., Price, V., McKereghan, K., Tushinski, R.J., Mochizuki, D.Y., Larsen, A., Grabstein, K., Gillis, S. and Cosman, D. *Proc. Natl. Acad. Sci. USA* 82: 6250-6254.
- Cerretti, D.P. et al. (1988) *Molecular Immunology* 25: 761.
- Chang, C.N., Rey, B., Bochner, B., Heyneker, H. and Gray, G. (1987) *Gene*: 189-196.
- 30 Cotes, P.M. and Bangham, D.R. (1961) *Nature* 191: 1065-1067
- Cox, G.N., McDermott, M.J., Merkel, E., Stroh, C.A., Ko, S.C., Squires, C.H., Gleason, T.M. and Russell, D. (1994) *Endocrinology* 135: 1913-1920.

- Cunningham, B.C. and Wells, J.A. (1989) *Science* 244: 1081-1085.
- Cunningham, B.C., Jhurani, P., Ng, P. and Wells, J.A. (1989) *Science* 243: 1330-1336.
- Cunningham, B.C., Ultsch, M., de Vos, A.M., Mulkerrin, M.G., Clauser, K.R. and Wells, J.A. (1991) *Science* 254: 821-825.
- 5 D'Andrea, A.D., Lodish, H.F. and Wong, G.G. (1989) *Cell* 57: 277-285.
- Davis, S., Aldrich, T.H., Stahl, N., Pan, L., Taga, T., Kishimoto, T., Ip, N.Y. and Yancopoulos, G.D. (1993) *Science* 260: 1805-1808.
- DeChiara, T.M., Erlitz, F. and Tarnowski, S.J. (1986) *Methods Enzymology*, 119: 403-15.
- de la Llosa, P., Chene, N. and Martal, J. (1985) *FEBS Letts.* 191: 211-215
- 10 Delorme, E., Lorenzini, T., Giffin, J., Martin, F., Jacobsen, F., Boone, T., Elliot, S. (1992) *Biochemistry* 31: 9871-9876.
- de Sauvage, F.J., Hass, P.E., Spencer, S.D. et al. (1994) *Nature* 369: 533-538.
- de Vos, A.M., Ultsch, M. and Kossiakoff, A.A. (1992) *Science* 255: 306-312.
- Devos, R., Plaetinck, G., Cheroutre, H., Simons, G., Degraeve, W., Tavernier, J., Remaut, E.
- 15 and Fiers, W. (1983) *Nucleic Acids Research* 11: 4307.
- Diederichs, K., Boone, T. and Karplus, A. (1991) *Science* 154: 1779-1782.
- Dorssers, L., Burger, H., Bot, F., Delwel, R., Van Kessel, A.H.M.G., Lowenberg, B. and Wagemaker, G. (1987) 55: 115-124.
- Dube, S., Fisher, J.W., Powell, J.S. (1988) *J. Biol. Chem.* 263: 17526-17521.
- 20 Fish, E.N., Banerjee, K., Levine, H.L. and Stebbing, N. (1986) *Antimicrob. Agents Chemother.* 30: 52-56.
- Foster, D.C., Sprecher, C.A., Grant, F.J., Kramer, J.M. et al. (1994) *Proc. Natl. Acad. Sci USA* 91: 13023-13027.
- Fukuda, M.N., Sasaki, H., Fukuda, M. (1989) *Blood* 73: 84-89.
- 25 Goeddel, D.V., Heyneker, H.L., Hozumi, T. et al., (1979) *Nature* 281: 544-548.
- Goeddel, D.V., Yelverton, E., Ullrich, A., Heyneker, H.L., Miozzari, G., Holmes, W., Seeburg, P.H., Dull, T., Mat, L., Stebbing, N., Crea, R., Maeda, S., McCandliss, R., Sloma, A., Tabor, J.M., Gross, M., Familletti, P.C. and Pestka, S. (1980) *Nature* 287: 411-416.
- Goldwasser, E. and Gross, M. (1975) *Methods Enzymology* 37: 109-121.
- 30 Goodson, R.J. and Katre, N.V. (1990) *Biotechnology* 8: 343-346.
- Goodwin, R.G., Lupton, S., Schmieder, A., Hjerrild, K.J., Jerzy, R., Clevenger, W., Gillis, S., Cosman, D. and Namen, A.E. (1989) *Proc. Natl. Acad. Sci. USA* 86: 302-306.

- Gough, N.M. et al., (1988) *Proc. Natl. Acad. Sci. USA* 85: 2623-2627.
- Gubler, U., Chua, A.O., Schoenhaut, D.S., Dwyer, C.M., McComas, W., Motyka, R., Nabavi, N., Wolitzky, A.G., Quinn, P.M., Familletti, P.C. and Gately, M.K. (1991) *Proc. Natl. Acad. Sci. USA* 88: 4143-4147.
- 5 Hershfield, M.S., Buckley, R.H., Greenberg, M.L. et al., (1987) *N.Engl. J. Medicine* 316: 589-596.
- Hill, C.P., Osslund, T.D. and Eisenberg, D. (1993) *Proc. Natl. Acad. Sci. USA* 90:5167-5171.
- Hoffman, R.C., Andersen, H., Walker, K., Krakover, J.D., Patel, S., Stamm, M.R. and Osborn, S.G. (1996) *Biochemistry* 35: 14849-14861.
- 10 Hsiung, H.M., Mayne, N.G. and Becker, G.W. (1986) *Biotechnology* 4: 991-995.
- Hercus, T.R., Bagley, C.J., Cambareri, B., Dottore, M., Woodcock, J.M., Vadas, M.A., Shannon, M.F., and Lopez, A.F. (1994) *Proc. Natl. Acad. Sci. USA* 91: 5838-5842.
- Hirano, T., Yasukawa, K., Harada, H., Taga, T., Watanabe, Y., Matsuda, T., Kashiwamura, S.-I., Nakajima, K., Koyama, K., Iwamatsu, A., Tsunasawa, S., Sakiyama, F., Matsui, H.,
- 15 Takahara, Y., Taniguchi, T. and Kishimoto, T. (1986) *Nature* 324: 73-76.
- Horisberger, M.A. and Di Marco, S. (1995) *Pharmac. Ther.* 66: 507-534.
- Innis, M.A., Gelfand, D.H., Sninsky, J.J. and White, T.J. (1990) *PCR Protocols: A Guide to Methods and Applications*. Academic Press, San Diego, CA.
- Jacobs, K., Shoemaker, C., Rudersdorf, R. et al., (1985) *Nature* 313: 806-810.
- 20 Karpusas, M., Nolte, M., Benton, C.B., Meier, W., Lipscomb, W.N. and Goelz, S. (1997) *Proc. Natl. Acad. Sci. USA* 94: 11813-11818.
- Katre, N.V. (1990) *J. Immunology* 144: 209-213.
- Katre, N.V., Knauf, M.J. and Laird, W.J. (1987) *Proc. Natl. Acad. Sci. USA* 84: 1487-1491.
- Kawasaki, E.S. et al., (1985) *Science* 230: 291.
- 25 Kawashima, I., Ohsumi, J., Mita-Honjo, K., Shimoda-Takano, Ishikawa, H., Sakakibara, S., Miyadai, K. and Takiguchi, Y. (1991) *FEBS Letts.* 283: 199-202.
- Kingsley, D.M. (1994) *Genes Dev.* 8: 133-146.
- Komatsu, N., Nakauchi, H., Miwa, A., Ishihara, T., Eguguchi, M. et al., (1991) *Cancer Research* 51, 341-348.
- 30 Kruse, N., Tony, H.-P. and Seibald, W. (1992) *EMBO J.* 11: 3237-3244.
- Lam, A., Fuller, F., Miller, J., Kloss, J., Manthorpe, M., Varon, S. and Cordell, B. (1991) *Gene* 102: 271-276.

- Kubota, N., Orita, T., Hattori, K., Oh-eda, M., Ochi, N. and Yamazaki, T. (1990) *J. Biochem.* 107: 486-492.
- Kuga, T., Komatsu, Y., Yamasaki, M., De4kine, S., Miyaji, H., Nishi, T., Sato, M., Yokoo, Y., Asano, M., Okabe, M., Morimoto, M. and Itoh, S. (1989) *Bioch.. Biophys. Res. Comm.* 159: 103-111.
- 5 Kunkel, T.A., Roberts, J.D. and Zakour, R.A. (1987) *Methods in Enzymology* 154: 367-382.
- Lange, B., Valtieri, M., Santoli, D., Caracciolo, D., Mavilio, F., Gemperlein, I., Griffin, C., Emanuel, B., Finan, J., Nowell, P. and Rovera, G. (1987) *Blood* 70: 192-199.
- Lee, F., Yokota, T., Otsuka, T., Gemmell, L., Larson, N., Luh, J., Arai, K.-I. and Rennick, D. 10 (1985) *Proc. Natl. Acad. Sci. USA* 82: 4360-4364.
- Lewis, J.A. (1995) Chapter 9: Antiviral activity of cytokines. pp. 129-141.
- Li, C.H. (1982) *Mol. Cell. Biochem.* 46: 31-41.
- Lin, F.-K. (1996) United States Patent # 5,547,933.
- Lin, F.-K., Suggs, S., Lin, C.-H., et al., *Proc. Natl. Acad. Sci. USA* (1985) 82: 7580-7584.
- 15 Linsley, P.S., Kallestad, J., Ochs, V. and Neubauer, M. (1990) 10: 1882-1890.
- Livnah, O., Stura, E.A., Johnson, D.L. et al., (1996) *Science* 273: 464-471.
- Lu, H.S., Clogston, C.L., Narhi, L.O., Merewether, L.A., Pearl, W.R. and Boone, T.C. (1992) *J. Biol. Chem.* 267: 8770-8777.
- Lydon, N.B., Favre, C., Bore, S., Neyret, O., Benureau, S., Levine, A.M., Seelig, G.F., 20 Nagabhushan, T. L. and Trotta, P.P. (1985) *Biochemistry* 24: 4131-41.
- MacGillivray, M.H., Baptista, J. and Johnson, A. (1996) *J. Clin. Endocrinol. Metab.* 81: 1806-1809.
- Malik, N., Kallestad, J.C., Gunderson, N.L., Austin, S.D., Neubauer, M.G., Ochs, V., Marquardt, H., Zarling, J.M., Shoyab, M., Wei, C.-M., Linsley, P.S. and Rose, T.M. (1989) 25 *Molecular and Cellular Biology* 9: 2847-2853.
- Martial, J., Chene, N. and de la Llosa, P. (1985) *FEBS. Letts.* 180: 295-299.
- Martial, J.A., Hallewell, R.A., Baxter, J.D. and Goodman, H.M. (1979) *Science* 205: 602-606.
- Matthews, D.J., Topping, R.S., Cass, R.T. and Giebel, L.B. (1996) *Proc. Natl. Acad. Sci.* 30 *USA* 93: 9471-9476.
- McKay, D.B. (1992) *Science* 257: 412.

- McKenzie, A.N.J., Culpepper, J.A., Malefyt, R. de W., Briere, F., Punnonen, J., Aversa, G., Sato, A., Dang, W., Cocks, B.G., Menon, S., de Vries, J.E., Banchereau, J., and Zurawski, G. (1993) *Proc. Natl. Acad. Sci. USA* 90: 3735-3739.
- Meyers, F.J., Paradise, C., Scudder, S.A., Goodman, G. and Konrad, M. (1991) *Clin. Pharmacol. Ther.* 49: 307-313.
- 5 Milburn, M.V., Hassell, A.M., Lambert, M.H., Jordan, S.R., Proudfoot, A.E., Graber, P. and Wells, T.N.C. (1993) *Nature* 363: 172-176.
- Mills, J.B., Kostyo, J.L., Reagan, C.R., Wagner, S.A., Moseley, M.H. and Wilhelm, A.E. (1980) *Endocrinology* 107: 391-399.
- 10 Minty, A., Chalon, P., Derocq, J.-M., Dumont, X., Guilemot, J.-C., Kaghad, M., Labit, C., Leplatois, P., Liauzun, P., Miloux, B., Minty, C., Casellas, P., Loison, G., Lupker, J., Shire, D., Ferrara, P. and Caput, D. (1993) *Nature* 362: 248-250.
- Moreau, J.-F., Donaldson, D.D., Bennett, F., Witek-Giannotti, J., Clark, S.C. and Wong, G.G. (1988) *Nature* 336: 690-692.
- 15 Mott, H.R. and Campbell, I.D. (1995) *Current Opinion in Structural Biology* 5: 114-121.
- Martial, J.A., Hallewell, R.A., Baxter, J.D. and Goodman, H.M. (1979) *Science* 205: 602-606.
- Nagata, S. (1994) in *Cytokines and Their Receptors*, N.A. Nicola ed., Oxford University Press, Oxford, pp. 158-160.
- 20 Nagata, S., Tsuchiya, M., Asano, S., Kziro, Y., Yamazaki, T., Yamamoto, O., Hirata, Y., Kubota, N., Oh-eda, M., Nomura, H. and Ono, M. (1986a) *Nature* 319: 415-418.
- Nagata, S., Tsuchiya, M., Asano, S., Yamamoto, O., Hirata, Y., Kubota, N., Oh-eda, M., Nomura, H. and Yamazaki, T. (1986b) *EMBO J.* 5: 575-581.
- O'Reilly, D.R., Miller, L.K., Luckow, V.A. (1992) *Caculovirus Expression Vectors*, W.H. Freeman Co. publishers, New York.
- 25 Otsuka, T., Miyajima, A., Brown, N., Otsu, K., Abrams, J., Saeland, S., Caux, C., De Waal-Malefyt, De Vries, J., Meyerson, P., Yokota, K., Gemmell, Rennick, D., Lee, F., Arai, N., Arai, K.-I. and Yokota, T. (1988) *J. Immunology* 140: 2288-2295.
- Owers-Narhi, L., Arakawa, T., Aoki, K.H., Elmore, R., Rohde, M.F., Boone, T., Strickland, T.W. (1991) *J. Biol. Chem.* 266: 23022-23026.
- 30 Ozes, O.S., Reiter, Z., Klein, S., Blatt, L.M. and Taylor, M.W. (1992) *J. Interferon Research* 12: 55-59.

- Paonessa, G., Graziani, R., de Serio, A., Savino, R., Ciapponi, L., Lahm, A., Ssalvati, A.L., Toniatti, C. and Ciliberto, G. (1995) *EMBO J.* 14: 1942-1951.
- Park, L.S., Waldron, P.E., Friend, D., Sassenfeld, H.M., Price, V., Anderson, D., Cosman, D., Andrews, R.G., Bernstein, I.D. and Urdal, D.L. (1989) *Blood* 74: 56-65.
- 5 Paul, S.R., Bennett, F., Calvetti, J.A., Kelleher, K., Wood, C.R., O'Hara, R.M., Leary, A.C., Sibley, B., Clark, S.C., William, D.A. and Yang, Y.-C. (1990) *Proc. Natl. Acad. Sci. USA* 87: 7512-7516.
- Pestka, S., Langer, J.A., Zoon, K.C. and Samuel, C.E. (1987) *Ann. Rev. Biochem.* 56: 727-777.
- 10 Pickering, L.A., Kronenberg, L.H. and Stewart, W.E., II (1980) *Proc. Natl. Acad. Sci. USA* 77: 5938-5942.
- Powers, R., Garrett, D.S., March, C.J., Frieden, E.A., Gronenborn, A.M. and Clore, G.M. (1992) *Science* 256:1673-1677.
- Radhakrishnan, R., Walter, L.J., Hruka, A., Reichert, P., Trotta, P.P., Nagabhushan, T. L. and
 15 Walter, M.R. (1996) *Structure* 4: 1453-1463.
- Read, L.C., Tomas, F.M., Howarth, G.S., Martin, A.A., Edson, K.J., Gillespie, C.M., Owens, P.C. and Ballard, F.J. (1992) *J. Endocrinology* 133: 421-431.
- Redfield, C., Smith L.J., Boyd, J., Lawrence, G.M.P., Edwards, R.G., Smith, R.A.G., and Dobson, C.M. (1991) *Biochemistry* 30: 11029-11035.
- 20 Robinson, R.C., Grey, L.M., Stauton, D., Vankelecom, H., Vernallis, A.B., Moreau, J.-F., Stuart, D.I., Heath, J.K. and Jones, E.Y. (1994) *Cell* 77: 1101-1116.
- Sasaki, H., Bothner, B., Dell, A. and Fukuda, M. (1987) *J. Biol. Chem.* 262: 12059-12076.
- Sasaki, H. Ochi, N., Dell, A. and Fukuda, M. (1988) *Biochemistry* 27: 8616-8626.
- Shaw, G., Veldman, G. and Wooters, J.L. (1992) United States Patent 5,166,322.
- 25 Shirafuji, N., Asano, S., Matsuda, S., Watari, K., Takaku, F. and Nagata, S. (1989) *Exp Hematol.* 17: 116-119.
- Silvennoinen, O. and Ihle, J.N. (1996) *Signalling by the Hematopoietic Cytokine Receptors*, R.G. Landes, Company, Austin, TX.
- Souza, L.M., Boone, T.C., Gabilove, J., Lai, P.H., Zsebo, K.M., Murdock, D.C., Chazin,
 30 V.R., Bruszewski, J., Lu, H., Chen, K.K., Barendt, J., Platzer, E., Moore, M.A.S., Mertelsmann, R. and Welte, K. (1986) *Science* 232: 61-65.
- Spivak, J.L., Hogans, B.B. (1989) *Blood* 73: 90-99.

- Takeuchi, M., Takasaki, S., Miyazaki, H., Takashi, D., Hoshi, S., Kochibe, N. and Kotaba, A. (1988) *J. Biol. Chem.* 263: 3657-3663.
- Tanaka, H., Satake-Ishikawa, R., Ishikawa, M., Matsuki, S. and Asano, K. (1991) *Cancer Research* 51: 3710-3714.
- 5 Taniguchi, T., Matsui, H., Fujita, T., Takaoka, C., Kashima, N., Yoshimoto, R. and Hamuro, J. (1983) *Nature* 302: 305-310.
- Taniguchi, T., Ohno, S., Fujii-Kuriyama, Y. and Muramatsu, M. (1980) *Gene* 10: 11-15.
- Tarnowski, S.J., Roy, S.K., Liptak, R.A., Lee, D.K. and Ning, R.Y. (1986) *Methods Enzymology* 119:153-165.
- 10 Tavernier, J., Tuypens, T., Verhee, A., Plaetinck, G., Devos, R., Van Der Heyden, J., Guisez, Y. and Oefner, C. (1995) *Proc. Natl. Acad. Sci. USA* 92: 5194-5198.
- Teh, L.-C. and Chapman, G.E. (1988) *Biochem. Biophys. Res. Comm.* 150: 391-398.
- Thatcher, D.R. and Panayotatos, N. (1986) *Methods Enzymology* 119: 166-177.
- Tomas, F.M., Knowles, S.E., Owens, P.C., Chandler, C.S., Francis, G.L., Read, L.C. and
- 15 Ballard, F.J. (1992) *Biochem. J.* 282: 91-97.
- Tsuchiya, M., Asano, S., Kaziro, Y and Nagata, S. (1986) *Proc. Natl. Acad. Sci. USA* 83: 7633-7637.
- Tsuda, E., Kawanishi, G., Ueda, M., Masuda, S. and Sasaki, R. (1990) *Eur. J. Biochem.* 188: 405-411.
- 20 Vieira, P., de Waal-Malefyt, R., Dang, M.N., Johnson, K.E., Kastelein, R., Fiorentino, D.F., de Vries, J.E., Roncarolo, M.-G., Mosmann, T.R. and Moore, K.W. (1991) *Proc. Natl. Acad. Sci. USA* 88: 1172-1176.
- Wang, A., Lu, S. -D., and Mark, D.F. (1984) *Science* 224: 1431-1433.
- Walter, M.R., Cook, W.J., Ealick, S.E., Nagabhusan, T.L., Trotta, P.T. and Bugg, C.E. (1992)
- 25 *J.Mol. Biol.* 224: 1075-1085.
- Wen, D., Boissel, J.-P. R., Tracy, T.E., Gruninger, R.H., Mulcahy, L.S., Czelusniak, J., Goodman, M. and Bunn, H.F. (1993) *Blood*: 1507-1516.
- Wen, D., Boissel, J.P., Showers, M., Ruch, B.C. and Bunn, H.F. (1994) *J. Biol. Chem.* 269: 22839-22846.
- 30 White, B. A., (1993), *Methods in Molecular Biology*, volume 15: PCR Protocols: Current Methods and Applications. Humana Press, Totowa, NJ.

- Wojchowski, D.M., Orkin, S.H., Sytkowshi, A.J. (1987) *Biochim. Biophys. Acta* 910: 224-232.
- Wolf, S.F., Temple, P.A., Kobayashi, M., Young, D., Dicig, M., Lowe, L., Dzialo, R., Fitz, L., Ferenz, C., Hewick, R.M., Kelleher, K., Herrmann, S.H., Clark, S.C., Azzoni, L.,
5 Chan, S.H., Trinchieri, G. and Perussia, B. (1991) *J. Immunology* 146: 3074-3081.
- Wong, G.G. et al., (1987) *Science* 235: 1504.
- Wrighton, N.C., Farrell, F.X., Chang, R. et al., (1996) *Science* 273: 458-463.
- Yamaguchi, K., Akai, K., Kawanishi, G. Ueda, M., Masuda, S., Sasaki, R. (1991). *J. Biol. Chem.* 266: 20434-20439.
- 10 Yang, Y.-C., Ciarletta, A.B., Temple, P.A., Chung, M.P., Kovacic, S., Witek-Giannotti, J.S., Leary, A.C., Kriz, R., Donahue, R.E., Wong, G.G. and Clark, S.C. (1986) *Cell* 47: 3-10.
- Yang, Y.-C., Ricciadi, S., Ciarletta, A., Calvetti, J., Kelleher, K. and Clark, S.C. (1989) *Blood* 74: 1880-1884.
- Yokota, T., Otsuka, T., Mosmann, T., Banchereau, J., DeFrance, T., Blanchard, D., De Vries, J.E., Lee F. and Arai, K.-I. (1986) *Proc. Natl. Acad. Sci. USA* 83: 5894-5898.
- 15 Yokota, T., Coffman, R.L., Hagiwara, H., Rennick, D.M., Takebe, Y., Yokota, K., Gemmell, L., Shrader, B., Yang, G., Meyerson, P., Luh, J., Hoy, P., Pene, J., Briere, F., Spits, H., Banchereau, J., De Vries, J., Lee, F., Arai, N. and Arai, K.-I. (1987) *Proc. Natl. Acad. Sci. USA* 84: 7388-7392.
- 20 Zurawski, S.M., Imler, J.L., and Zurawski, G. (1990) *EMBO J.* 9: 3899-3905.
- Zurawski, S.M. and Zurawski, G. (1992) *EMBO J.* 11: 3905-3910.

What is claimed is:

1. A cysteine variant of a member of the GH supergene family comprising a cysteine residue substituted for an amino acid selected from the group consisting of an amino acid in the loop regions, an amino acid near the ends of the alpha helices, an amino acid proximal to the first amphipathic helix, and an amino acid distal to the final amphipathic helix or wherein the cysteine residue is added at the N-terminus or C-terminus of the proteins.
2. A cysteine variant of a member of the GH supergene family comprising a cysteine residue introduced between two amino acids in the loop regions, the ends of the alpha helices, proximal to the first amphipathic helix, or distal to the final amphipathic helix.
3. A cysteine variant according to claim 1 wherein the amino acid substituted for is part of an N- or O-linked glycosylation site.
4. A cysteine variant according to claim 2 wherein the cysteine residue is introduced between two amino acids in an N-linked glycosylation site or adjacent to an amino acid in an N-linked or O-linked glycosylation site.
5. A cysteine variant according to claims 1-4 wherein the member of the GH supergene family is selected from the group consisting of growth hormone, prolactin, placental lactogen, erythropoietin, thrombopoietin, interleukin-2, interleukin-3, interleukin-4, interleukin-5, interleukin-6, interleukin-7, interleukin-9, interleukin-10, interleukin-11, interleukin-12 (p35 subunit), interleukin-13, interleukin-15, oncostatin M, ciliary neurotrophic factor, leukemia inhibitory factor, alpha interferon, beta interferon, gamma interferon, omega interferon, tau interferon, granulocyte-colony stimulating factor, granulocyte-macrophage colony stimulating factor, cardiotrophin-1 and macrophage colony stimulating factor.
6. A cysteine variant according to claim 1 wherein the amino acid substituted for is in the A-B loop, the B-C loop, the C-D loop or D-E loop of interferon/interferon-10-like members of the GH supergene family.

7. A cysteine variant according to claim 2 wherein the loop region is the A-B loop, the B-C loop, the C-D loop or D-E loop of interferon/interferon-10-like members of the GH supergene family.
- 5 8. A cysteine variant according to claims 1-7 wherein the added cysteine is PEGylated.
9. A cysteine variant according to claim 1 wherein the member of the GH supergene family is growth hormone.
- 10 10. The cysteine variant according to claim 9 wherein the substituted for amino acid is selected from the group consisting of amino acids located at the N-terminal end of the A-B loop, the B-C loop, the C-D loop, the first three or last three amino acids in the A, B, C and D helices and the amino acids proximal to helix A and distal to helix D.
- 15 11. The cysteine variant according to claim 10 wherein the substituted for amino acid is selected from the group consisting of F1, T3, P5, E33, A34, K38, E39, Q40, S43, Q46, N47, P48, Q49, T50, S51, S55, T60, A98, N99, S100, G104, A105, S106, E129, D130, G131, S132, P133, T135, G136, Q137, K140, Q141, T142, S144, K145, D147, T148, N149, S150, H151, N152, D153, S184, E186, G187, S188, and G190.
- 20 12. A cysteine variant according to claim 1 wherein the member of the GH supergene family is erythropoietin.
13. The cysteine variant according to claim 12 wherein the substituted for amino acid is
25 selected from the group consisting of amino acids located in the A-B loop, the B-C loop, the C-D loop, the amino acids proximal to helix A and distal to helix D and the N- or C-terminus.
14. A cysteine variant according to claim 13 wherein the substituted for amino acid is selected
30 from the group consisting of serine-126, N24, I25, T26, N38, I39, T40, N83, S84, A1, P2, P3, R4, D8, S9, T27, G28, A30, E31, H32, S34, N36, D43, T44, K45, N47, A50, K52, E55, G57, Q58, G77, Q78, A79, Q86, W88, E89, T107, R110, A111, G113, A114, Q115,

K116, E117, A118, S120, P121, P122, D123, A124, A125, A127, A128, T132, K154, T157, G158, E159, A160, T163, G164, D165, R166 and S85.

- 5 **15.** A cysteine variant according to claim 2 wherein the member of the GH supergene family is growth hormone.
- 10 **16.** A cysteine variant according to claim 15 wherein the cysteine is introduced into the region selected from the group consisting of amino acids located at the N-terminal end of the A-B loop, the B-C loop, the C-D loop, the first three or last three amino acids in the A, B, C and D helices and the amino acids proximal to helix A and distal to helix D.
- 15 **17.** A cysteine variant according to claim 2 wherein the member of the GH supergene family is erythropoietin.
- 15 **18.** A cysteine variant according to claim 17 wherein the cysteine is introduced into the region selected from the group consisting of the N-terminal end of the A-B loop, the B-C loop, the C-D loop, the area of the first three or last three amino acids in the A, B, C and D helices and proximal to helix A and distal to helix D.
- 20 **19.** A cysteine variant according to claim 2 wherein the member of the GH supergene family is alpha interferon.
- 25 **20.** A cysteine variant according to claim 19 wherein the cysteine is introduced into the region selected from the group consisting of the N-terminal end of the A-B loop, the B-C loop, the C-D loop, the area of the first three or last three amino acids in the A, B, C and D helices and proximal to helix A and distal to helix D.
- 30 **21.** A cysteine variant according to claim 2 wherein the member of the GH supergene family is granulocyte-colony stimulating factor.
- 22.** A cysteine variant according to claim 21 wherein the cysteine is introduced into the region selected from the group consisting of the N-terminal end of the A-B loop, the B-C loop,

the C-D loop, the area of the first three or last three amino acids in the A, B, C and D helices and proximal to helix A and distal to helix D.

23. A cysteine variant according to claims 8-22 wherein the added cysteine is PEGylated.

SEQUENCE LISTING

<110> Cox III, George N
Bolder Biotechnology, Inc.

<120> Derivatives of Growth Hormone and Related Proteins

<130> BB0011

<140> unknown

<141> 1998-07-14

<150> 60/052,516

<151> 1997-07-14

<160> 41

<170> PatentIn Ver. 2.0

<210> 1

<211> 191

<212> PRT

<213> Homo sapiens

<400> 1

Phe Pro Thr Ile Pro Leu Ser Arg Leu Phe Asp Asn Ala Met Leu Arg
1 5 10 15

Ala His Arg Leu His Gln Leu Ala Phe Asp Thr Tyr Gln Glu Phe Glu
20 25 30

Glu Ala Tyr Ile Pro Lys Glu Gln Lys Tyr Ser Phe Leu Gln Asn Pro
35 40 45

Gln Thr Ser Leu Cys Phe Ser Glu Ser Ile Pro Thr Pro Ser Asn Arg
50 55 60

Glu Glu Thr Gln Gln Lys Ser Asn Leu Glu Leu Leu Arg Ile Ser Leu
65 70 75 80

Leu Leu Ile Gln Ser Trp Leu Glu Pro Val Gln Phe Leu Arg Ser Val
85 90 95

Phe Ala Asn Ser Leu Val Tyr Gly Ala Ser Asp Ser Asn Val Tyr Asp
100 105 110

Leu Leu Lys Asp Leu Glu Glu Gly Ile Gln Thr Leu Met Gly Arg Leu
115 120 125

Glu Asp Gly Ser Pro Arg Thr Gly Gln Ile Phe Lys Gln Thr Tyr Ser
130 135 140

Lys Phe Asp Thr Asn Ser His Asn Asp Asp Ala Leu Leu Lys Asn Tyr
145 150 155 160

Gly Leu Leu Tyr Cys Phe Arg Lys Asp Met Asp Lys Val Glu Thr Phe
165 170 175

Leu Arg Ile Val Gln Cys Arg Ser Val Glu Gly Ser Cys Gly Phe
180 185 190

<210> 2
 <211> 166
 <212> PRT
 <213> Homo sapiens

<400> 2
 Ala Pro Pro Arg Leu Ile Cys Asp Ser Arg Val Leu Glu Arg Tyr Leu
 1 5 10 15
 Leu Glu Ala Lys Glu Ala Glu Asn Ile Thr Thr Gly Cys Ala Glu His
 20 25 30
 Cys Ser Leu Asn Glu Asn Ile Thr Val Pro Asp Thr Lys Val Asn Phe
 35 40 45
 Tyr Ala Trp Lys Arg Met Glu Val Gly Gln Gln Ala Val Glu Val Trp
 50 55 60
 Gln Gly Leu Ala Leu Leu Ser Glu Ala Val Leu Arg Gly Gln Ala Leu
 65 70 75 80
 Leu Val Asn Ser Ser Gln Pro Trp Glu Pro Leu Gln Leu His Val Asp
 85 90 95
 Lys Ala Val Ser Gly Leu Arg Ser Leu Thr Thr Leu Leu Arg Ala Leu
 100 105 110
 Gly Ala Gln Lys Glu Ala Ile Ser Pro Pro Asp Ala Ala Ser Ala Ala
 115 120 125
 Pro Leu Arg Thr Ile Thr Ala Asp Thr Phe Arg Lys Leu Phe Arg Val
 130 135 140
 Tyr Ser Asn Phe Leu Arg Gly Lys Leu Lys Leu Tyr Thr Gly Glu Ala
 145 150 155 160
 Cys Arg Thr Gly Asp Arg
 165

<210> 3
 <211> 165
 <212> PRT
 <213> Homo sapiens

<400> 3
 Cys Asp Leu Pro Gln Thr His Ser Leu Gly Ser Arg Arg Thr Leu Met
 1 5 10 15
 Leu Leu Ala Gln Met Arg Arg Ile Ser Leu Phe Ser Cys Leu Lys Asp
 20 25 30
 Arg His Asp Phe Gly Phe Pro Gln Glu Glu Phe Gly Asn Gln Phe Gln
 35 40 45
 Lys Ala Glu Thr Ile Pro Val Leu His Glu Met Ile Gln Gln Ile Phe
 50 55 60
 Asn Leu Phe Ser Thr Lys Asp Ser Ser Ala Ala Trp Asp Glu Thr Leu
 65 70 75 80

Leu Asp Lys Phe Tyr Thr Glu Leu Tyr Gln Gln Leu Asn Asp Leu Glu
 85 90 95
 Ala Cys Val Ile Gln Gly Val Gly Val Thr Glu Thr Pro Leu Met Lys
 100 105 110
 Glu Asp Ser Ile Leu Ala Val Arg Lys Tyr Phe Gln Arg Ile Thr Leu
 115 120 125
 Tyr Leu Lys Glu Lys Lys Tyr Ser Pro Cys Ala Trp Glu Val Val Arg
 130 135 140
 Ala Glu Ile Met Arg Ser Phe Ser Leu Ser Thr Asn Leu Gln Glu Ser
 145 150 155 160
 Leu Arg Ser Lys Glu
 165

<210> 4
 <211> 166
 <212> PRT
 <213> Homo sapiens

<400> 4
 Cys Asp Leu Pro Glu Thr His Ser Leu Asp Asn Arg Arg Thr Leu Met
 1 5 10 15
 Leu Leu Ala Gln Met Ser Arg Ile Ser Pro Ser Ser Cys Leu Met Asp
 20 25 30
 Arg His Asp Phe Gly Phe Pro Gln Glu Glu Phe Asp Gly Asn Gln Phe
 35 40 45
 Gln Lys Ala Pro Ala Ile Ser Val Leu His Glu Leu Ile Gln Gln Ile
 50 55 60
 Phe Asn Leu Phe Thr Thr Lys Asp Ser Ser Ala Ala Trp Asp Glu Asp
 65 70 75 80
 Leu Leu Asp Lys Phe Cys Thr Glu Leu Tyr Gln Gln Leu Asn Asp Leu
 85 90 95
 Glu Ala Cys Val Met Gln Glu Glu Arg Val Gly Glu Thr Pro Leu Met
 100 105 110
 Asn Ala Asp Ser Ile Leu Ala Val Lys Lys Tyr Phe Arg Arg Ile Thr
 115 120 125
 Leu Tyr Leu Thr Glu Lys Lys Tyr Ser Pro Cys Ala Trp Glu Val Val
 130 135 140
 Arg Ala Glu Ile Met Arg Ser Leu Ser Leu Ser Thr Asn Leu Gln Glu
 145 150 155 160
 Arg Leu Arg Arg Lys Glu
 165

<210> 5

<211> 166
 <212> PRT
 <213> Homo sapiens

<400> 5

```

Met Ser Tyr Asn Leu Leu Gly Phe Leu Gln Arg Ser Ser Asn Phe Gln
 1              5              10              15

Cys Gln Lys Leu Leu Trp Gln Leu Asn Gly Arg Leu Glu Tyr Cys Leu
      20              25              30

Lys Asp Arg Met Asn Phe Asp Ile Pro Glu Glu Ile Lys Gln Leu Gln
      35              40              45

Gln Phe Gln Lys Glu Asp Ala Ala Leu Thr Ile Tyr Glu Met Leu Gln
      50              55              60

Asn Ile Phe Ala Ile Phe Arg Gln Asp Ser Ser Ser Thr Gly Trp Asn
      65              70              75              80

Glu Thr Ile Val Glu Asn Leu Leu Ala Asn Val Tyr His Gln Ile Asn
      85              90              95

His Leu Lys Thr Val Leu Glu Glu Lys Leu Glu Lys Glu Asp Phe Thr
      100             105             110

Arg Gly Lys Leu Met Ser Ser Leu His Leu Lys Arg Tyr Tyr Gly Arg
      115             120             125

Ile Leu His Tyr Leu Lys Ala Lys Glu Tyr Ser His Cys Ala Trp Thr
      130             135             140

Ile Val Arg Val Glu Ile Leu Arg Asn Phe Tyr Phe Ile Asn Arg Leu
      145             150             155             160

Thr Gly Tyr Leu Arg Asn
      165
  
```

<210> 6
 <211> 174
 <212> PRT
 <213> Homo sapiens

<400> 6

```

Thr Pro Leu Gly Pro Ala Ser Ser Leu Pro Gln Ser Phe Leu Leu Lys
 1              5              10              15

Cys Leu Glu Gln Val Arg Lys Ile Gln Gly Asp Gly Ala Ala Leu Gln
      20              25              30

Glu Lys Leu Cys Ala Thr Tyr Lys Leu Cys His Pro Glu Glu Leu Val
      35              40              45

Leu Leu Gly His Ser Leu Gly Ile Pro Trp Ala Pro Leu Ser Ser Cys
      50              55              60

Pro Ser Gln Ala Leu Gln Leu Ala Gly Cys Leu Ser Gln Leu His Ser
      65              70              75              80
  
```

Gly Leu Phe Leu Tyr Gln Gly Leu Leu Gln Ala Leu Glu Gly Ile Ser
 85 90 95
 Pro Glu Leu Gly Pro Thr Leu Asp Thr Leu Gln Leu Asp Val Ala Asp
 100 105 110
 Phe Ala Thr Thr Ile Trp Gln Gln Met Glu Glu Leu Gly Met Ala Pro
 115 120 125
 Ala Leu Gln Pro Thr Gln Gly Ala Met Pro Ala Phe Ala Ser Ala Phe
 130 135 140
 Gln Arg Arg Ala Gly Gly Val Leu Val Ala Ser His Leu Gln Ser Phe
 145 150 155 160
 Leu Glu Val Ser Tyr Arg Val Leu Arg His Leu Ala Gln Pro
 165 170

<210> 7
 <211> 332
 <212> PRT
 <213> Homo sapiens

<400> 7
 Ser Pro Ala Pro Pro Ala Cys Asp Leu Arg Val Leu Ser Lys Leu Leu
 1 5 10 15
 Arg Asp Ser His Val Leu His Ser Arg Leu Ser Gln Cys Pro Glu Val
 20 25 30
 His Pro Leu Pro Thr Pro Val Leu Leu Pro Ala Val Asp Phe Ser Leu
 35 40 45
 Gly Glu Trp Lys Thr Gln Met Glu Glu Thr Lys Ala Gln Asp Ile Leu
 50 55 60
 Gly Ala Val Thr Leu Leu Leu Glu Gly Val Met Ala Ala Arg Gly Gln
 65 70 75 80
 Leu Gly Pro Thr Cys Leu Ser Ser Leu Leu Gly Gln Leu Ser Gly Gln
 85 90 95
 Val Arg Leu Leu Leu Gly Ala Leu Gln Ser Leu Leu Gly Thr Gln Leu
 100 105 110
 Pro Pro Gln Gly Arg Thr Thr Ala His Lys Asp Pro Asn Ala Ile Phe
 115 120 125
 Leu Ser Phe Gln His Leu Leu Arg Gly Lys Val Arg Phe Leu Met Leu
 130 135 140
 Val Gly Gly Ser Thr Leu Cys Val Arg Arg Ala Pro Pro Thr Thr Ala
 145 150 155 160
 Val Pro Ser Arg Thr Ser Leu Val Leu Thr Leu Asn Glu Leu Pro Asn
 165 170 175
 Arg Thr Ser Gly Leu Leu Glu Thr Asn Phe Thr Ala Ser Ala Arg Thr
 180 185 190

Thr Gly Ser Gly Leu Leu Lys Trp Gln Gln Gly Phe Arg Ala Lys Ile
 195 200 205
 Pro Gly Leu Leu Asn Gln Thr Ser Arg Ser Leu Asp Gln Ile Pro Gly
 210 215 220
 Tyr Leu Asn Arg Ile His Glu Leu Leu Asn Gly Thr Arg Gly Leu Phe
 225 230 235 240
 Pro Gly Pro Ser Arg Arg Thr Leu Gly Ala Pro Asp Ile Ser Ser Gly
 245 250 255
 Thr Ser Asp Thr Gly Ser Leu Pro Pro Asn Leu Gln Pro Gly Tyr Ser
 260 265 270
 Pro Ser Pro Thr His Pro Pro Thr Gly Gly Tyr Thr Leu Phe Pro Leu
 275 280 285
 Pro Pro Thr Leu Pro Thr Pro Val Val Gln Leu His Pro Leu Leu Pro
 290 295 300
 Asp Pro Ser Ala Pro Thr Pro Thr Pro Thr Ser Pro Leu Leu Asn Thr
 305 310 315 320
 Ser Tyr Thr His Ser Gln Asn Leu Ser Gln Glu Gly
 325 330

<210> 8

<211> 127

<212> PRT

<213> Homo sapiens

<400> 8

Ala Pro Ala Arg Ser Pro Ser Pro Ser Thr Gln Pro Trp Glu His Val
 1 5 10 15
 Asn Ala Ile Gln Glu Ala Arg Arg Leu Leu Asn Leu Ser Arg Asp Thr
 20 25 30
 Ala Ala Glu Met Asn Glu Thr Val Glu Val Ile Ser Glu Met Phe Asp
 35 40 45
 Leu Gln Glu Pro Thr Cys Leu Gln Thr Arg Leu Glu Leu Tyr Lys Gln
 50 55 60
 Gly Leu Arg Gly Ser Leu Thr Lys Leu Lys Gly Pro Leu Thr Met Met
 65 70 75 80
 Ala Ser His Tyr Lys Gln His Cys Pro Pro Thr Pro Glu Thr Ser Cys
 85 90 95
 Ala Thr Gln Ile Ile Thr Phe Glu Ser Phe Lys Glu Asn Leu Lys Asp
 100 105 110
 Phe Leu Leu Val Ile Pro Phe Asp Cys Trp Glu Pro Val Gln Glu
 115 120 125

<210> 9

<211> 133
 <212> PRT
 <213> Homo sapiens

<400> 9

Ala Pro Thr Ser Ser Ser Thr Lys Lys Thr Gln Leu Gln Leu Glu His
 1 5 10 15
 Leu Leu Leu Asp Leu Gln Met Ile Leu Asn Gly Ile Asn Asn Tyr Lys
 20 25 30
 Asn Pro Lys Leu Thr Arg Met Leu Thr Phe Lys Phe Tyr Met Pro Lys
 35 40 45
 Lys Ala Thr Glu Leu Lys His Leu Gln Cys Leu Glu Glu Glu Leu Lys
 50 55 60
 Pro Leu Glu Glu Val Leu Asn Leu Ala Gln Ser Lys Asn Phe His Leu
 65 70 75 80
 Arg Pro Arg Asp Leu Ile Ser Asn Ile Asn Val Ile Val Leu Glu Leu
 85 90 95
 Lys Gly Ser Glu Thr Thr Phe Met Cys Glu Tyr Ala Asp Glu Thr Ala
 100 105 110
 Thr Ile Val Glu Phe Leu Asn Arg Trp Ile Thr Phe Cys Gln Ser Ile
 115 120 125
 Ile Ser Thr Leu Thr
 130

<210> 10
 <211> 152
 <212> PRT
 <213> Homo sapiens

<400> 10

Met Ser Arg Leu Pro Val Leu Leu Leu Leu Gln Leu Leu Val Arg Pro
 1 5 10 15
 Gly Leu Gln Ala Pro Met Thr Gln Thr Thr Pro Leu Lys Thr Ser Trp
 20 25 30
 Val Asn Cys Ser Asn Met Ile Asp Glu Ile Ile Thr His Leu Lys Gln
 35 40 45
 Pro Pro Leu Pro Leu Leu Asp Phe Asn Asn Leu Asn Gly Glu Asp Gln
 50 55 60
 Asp Ile Leu Met Glu Asn Asn Leu Arg Arg Pro Asn Leu Glu Ala Phe
 65 70 75 80
 Asn Arg Ala Val Lys Ser Leu Gln Asn Ala Ser Ala Ile Glu Ser Ile
 85 90 95
 Leu Lys Asn Leu Leu Pro Cys Leu Pro Leu Ala Thr Ala Ala Pro Thr
 100 105 110
 Arg His Pro Ile His Ile Lys Asp Gly Asp Trp Asn Glu Phe Arg Arg

115 120 125
 Lys Leu Thr Phe Tyr Leu Lys Thr Leu Glu Asn Ala Gln Ala Gln Gln
 130 135 140
 Thr Thr Leu Ser Leu Ala Ile Phe
 145 150

<210> 11
 <211> 129
 <212> PRT
 <213> Homo sapiens

<400> 11
 His Lys Cys Asp Ile Thr Leu Gln Glu Ile Ile Lys Thr Leu Asn Ser
 1 5 10 15
 Leu Thr Glu Gln Lys Thr Leu Cys Thr Glu Leu Thr Val Thr Asp Ile
 20 25 30
 Phe Ala Ala Ser Lys Asn Thr Thr Glu Lys Glu Thr Phe Cys Arg Ala
 35 40 45
 Ala Thr Val Leu Arg Gln Phe Tyr Ser His His Glu Lys Asp Thr Arg
 50 55 60
 Cys Leu Gly Ala Thr Ala Gln Gln Phe His Arg His Lys Gln Leu Ile
 65 70 75 80
 Arg Phe Leu Lys Arg Leu Asp Arg Asn Leu Trp Gly Leu Ala Gly Leu
 85 90 95
 Asn Ser Cys Pro Val Lys Glu Ala Asn Gln Ser Thr Leu Glu Asn Phe
 100 105 110
 Leu Glu Arg Leu Lys Thr Ile Met Arg Glu Lys Tyr Ser Lys Cys Ser
 115 120 125
 Ser

<210> 12
 <211> 134
 <212> PRT
 <213> Homo sapiens

<400> 12
 Met Arg Met Leu Leu His Leu Ser Leu Leu Ala Leu Gly Ala Ala Tyr
 1 5 10 15
 Val Tyr Ala Ile Pro Thr Glu Ile Pro Thr Ser Ala Leu Val Lys Glu
 20 25 30
 Thr Leu Ala Leu Leu Ser Thr His Arg Thr Leu Leu Ile Ala Asn Glu
 35 40 45
 Thr Leu Arg Ile Pro Val Pro Val His Lys Asn His Gln Leu Cys Thr
 50 55 60

Glu Glu Ile Phe Gln Gly Ile Gly Thr Leu Glu Ser Gln Thr Val Gln
 65 70 75 80
 Gly Gly Thr Val Glu Arg Leu Phe Lys Asn Leu Ser Leu Ile Lys Lys
 85 90 95
 Tyr Ile Asp Gly Gln Lys Lys Lys Cys Gly Glu Glu Arg Arg Arg Val
 100 105 110
 Asn Gln Phe Leu Asp Tyr Leu Gln Glu Phe Leu Gly Val Met Asn Thr
 115 120 125
 Glu Trp Ile Ile Glu Ser
 130

<210> 13
 <211> 212
 <212> PRT
 <213> Homo sapiens

<400> 13
 Met Asn Ser Phe Ser Thr Ser Ala Phe Gly Pro Val Ala Phe Ser Leu
 1 5 10 15
 Gly Leu Leu Leu Val Leu Pro Ala Ala Phe Pro Ala Pro Val Pro Pro
 20 25 30
 Gly Glu Asp Ser Lys Asp Val Ala Ala Pro His Arg Gln Pro Leu Thr
 35 40 45
 Ser Ser Glu Arg Ile Asp Lys Gln Ile Arg Tyr Ile Leu Asp Gly Ile
 50 55 60
 Ser Ala Leu Arg Lys Glu Thr Cys Asn Lys Ser Asn Met Cys Glu Ser
 65 70 75 80
 Ser Lys Glu Ala Leu Ala Glu Asn Asn Leu Asn Leu Pro Lys Met Ala
 85 90 95
 Glu Lys Asp Gly Cys Phe Gln Ser Gly Phe Asn Glu Glu Thr Cys Leu
 100 105 110
 Val Lys Ile Ile Thr Gly Leu Leu Glu Phe Glu Val Tyr Leu Glu Tyr
 115 120 125
 Leu Gln Asn Arg Phe Glu Ser Ser Glu Glu Gln Ala Arg Ala Val Gln
 130 135 140
 Met Ser Thr Lys Val Leu Ile Gln Phe Leu Gln Lys Lys Ala Lys Asn
 145 150 155 160
 Leu Asp Ala Ile Thr Thr Pro Asp Pro Thr Thr Asn Ala Ser Leu Leu
 165 170 175
 Thr Lys Leu Gln Ala Gln Asn Gln Trp Leu Gln Asp Met Thr Thr His
 180 185 190

Leu Ile Leu Arg Ser Phe Lys Glu Phe Leu Gln Ser Ser Leu Arg Ala
 195 200 205

Leu Arg Gln Met
 210

<210> 14
 <211> 177
 <212> PRT
 <213> Homo sapiens

<400> 14
 Met Phe His Val Ser Phe Arg Tyr Ile Phe Gly Leu Pro Pro Leu Ile
 1 5 10 15

Leu Val Leu Leu Pro Val Ala Ser Ser Asp Cys Asp Ile Glu Gly Lys
 20 25 30

Asp Gly Lys Gln Tyr Glu Ser Val Leu Met Val Ser Ile Asp Gln Leu
 35 40 45

Leu Asp Ser Met Lys Glu Ile Gly Ser Asn Cys Leu Asn Asn Glu Phe
 50 55 60

Asn Phe Phe Lys Arg His Ile Cys Asp Ala Asn Lys Glu Gly Met Phe
 65 70 75 80

Leu Phe Arg Ala Ala Arg Lys Leu Arg Gln Phe Leu Lys Met Asn Ser
 85 90 95

Thr Gly Asp Phe Asp Leu His Leu Leu Lys Val Ser Glu Gly Thr Thr
 100 105 110

Ile Leu Leu Asn Cys Thr Gly Gln Val Lys Gly Arg Lys Pro Ala Ala
 115 120 125

Leu Gly Glu Ala Gln Pro Thr Lys Ser Leu Glu Glu Asn Lys Ser Leu
 130 135 140

Lys Glu Gln Lys Lys Leu Asn Asp Leu Cys Phe Leu Lys Arg Leu Leu
 145 150 155 160

Gln Glu Ile Lys Thr Cys Trp Asn Lys Ile Leu Met Gly Thr Lys Glu
 165 170 175

His

<210> 15
 <211> 144
 <212> PRT
 <213> Homo sapiens

<400> 15
 Met Leu Leu Ala Met Val Leu Thr Ser Ala Leu Leu Leu Cys Ser Val
 1 5 10 15

Ala Gly Gln Gly Cys Pro Thr Leu Ala Gly Ile Leu Asp Ile Asn Phe
 20 25 30

Leu Ile Asn Lys Met Gln Glu Asp Pro Ala Ser Lys Cys His Cys Ser
 35 40 45
 Ala Asn Val Thr Ser Cys Leu Cys Leu Gly Ile Pro Ser Asp Asn Cys
 50 55 60
 Thr Arg Pro Cys Phe Ser Glu Arg Leu Ser Gln Met Thr Asn Thr Thr
 65 70 75 80
 Met Gln Thr Arg Tyr Pro Leu Ile Phe Ser Arg Val Lys Lys Ser Val
 85 90 95
 Glu Val Leu Lys Asn Asn Lys Cys Pro Tyr Phe Ser Cys Glu Gln Pro
 100 105 110
 Cys Asn Gln Thr Thr Ala Gly Asn Ala Leu Thr Phe Leu Lys Ser Leu
 115 120 125
 Leu Glu Ile Phe Gln Lys Glu Lys Met Arg Gly Met Arg Gly Lys Ile
 130 135 140

<210> 16
 <211> 178
 <212> PRT
 <213> Homo sapiens

<400> 16
 Met His Ser Ser Ala Leu Leu Cys Cys Leu Val Leu Leu Thr Gly Val
 1 5 10 15
 Arg Ala Ser Pro Gly Gln Gly Thr Gln Ser Glu Asn Ser Cys Thr His
 20 25 30
 Phe Pro Gly Asn Leu Pro Asn Met Leu Arg Asp Leu Arg Asp Ala Phe
 35 40 45
 Ser Arg Val Lys Thr Phe Phe Gln Met Lys Asp Gln Leu Asp Asn Leu
 50 55 60
 Leu Leu Lys Glu Ser Leu Leu Glu Asp Phe Lys Gly Tyr Leu Gly Cys
 65 70 75 80
 Gln Ala Leu Ser Glu Met Ile Gln Phe Tyr Leu Glu Glu Val Met Pro
 85 90 95
 Gln Ala Glu Asn Gln Asp Pro Asp Ile Lys Ala His Val Asn Ser Leu
 100 105 110
 Gly Glu Asn Leu Lys Thr Leu Arg Leu Arg Leu Arg Arg Cys His Arg
 115 120 125
 Phe Leu Pro Cys Glu Asn Lys Ser Lys Ala Val Glu Gln Val Lys Asn
 130 135 140
 Ala Phe Asn Lys Leu Gln Glu Lys Gly Ile Tyr Lys Ala Met Ser Glu
 145 150 155 160

Phe Asp Ile Phe Ile Asn Tyr Ile Glu Ala Tyr Met Thr Met Lys Ile
 165 170 175

Arg Asn

<210> 17
 <211> 199
 <212> PRT
 <213> Homo sapiens

<400> 17
 Met Asn Cys Val Cys Arg Leu Val Leu Val Val Leu Ser Leu Trp Pro
 1 5 10 15
 Asp Thr Ala Val Ala Pro Gly Pro Pro Pro Gly Pro Pro Arg Val Ser
 20 25 30
 Pro Asp Pro Arg Ala Glu Leu Asp Ser Thr Val Leu Leu Thr Arg Ser
 35 40 45
 Leu Leu Ala Asp Thr Arg Gln Leu Ala Ala Gln Leu Arg Asp Lys Phe
 50 55 60
 Pro Ala Asp Gly Asp His Asn Leu Asp Ser Leu Pro Thr Leu Ala Met
 65 70 75 80
 Ser Ala Gly Ala Leu Gly Ala Leu Gln Leu Pro Gly Val Leu Thr Arg
 85 90 95
 Leu Arg Ala Asp Leu Leu Ser Tyr Leu Arg His Val Gln Trp Leu Arg
 100 105 110
 Arg Ala Gly Gly Ser Ser Leu Lys Thr Leu Glu Pro Glu Leu Gly Thr
 115 120 125
 Leu Gln Ala Arg Leu Asp Arg Leu Leu Arg Arg Leu Gln Leu Leu Met
 130 135 140
 Ser Arg Leu Ala Leu Pro Gln Pro Pro Pro Asp Pro Pro Ala Pro Pro
 145 150 155 160
 Leu Ala Pro Pro Ser Ser Ala Trp Gly Gly Ile Arg Ala Ala His Ala
 165 170 175
 Ile Leu Gly Gly Leu His Leu Thr Leu Asp Trp Ala Val Arg Gly Leu
 180 185 190
 Leu Leu Leu Lys Thr Arg Leu
 195

<210> 18
 <211> 219
 <212> PRT
 <213> Homo sapiens

<400> 18

```

Met Cys Pro Ala Arg Ser Leu Leu Leu Val Ala Thr Leu Val Leu Leu
 1              5              10              15

Asp His Leu Ser Leu Ala Arg Asn Leu Pro Val Ala Thr Pro Asp Pro
      20              25              30

Gly Met Phe Pro Cys Leu His His Ser Gln Asn Leu Leu Arg Ala Val
      35              40              45

Ser Asn Met Leu Gln Lys Ala Arg Gln Thr Leu Glu Phe Tyr Pro Cys
      50              55              60

Thr Ser Glu Glu Ile Asp His Glu Asp Ile Thr Lys Asp Lys Thr Ser
      65              70              75              80

Thr Val Glu Ala Cys Leu Pro Leu Glu Leu Thr Lys Asn Glu Ser Cys
      85              90              95

Leu Asn Ser Arg Glu Thr Ser Phe Ile Thr Asn Gly Ser Cys Leu Ala
      100              105              110

Ser Arg Lys Thr Ser Phe Met Met Ala Leu Cys Leu Ser Ser Ile Tyr
      115              120              125

Glu Asp Leu Lys Met Tyr Gln Val Glu Phe Lys Thr Met Asn Ala Lys
      130              135              140

Leu Leu Met Asp Pro Lys Arg Gln Ile Phe Leu Asp Gln Asn Met Leu
      145              150              155              160

Ala Val Ile Asp Glu Leu Met Gln Ala Leu Asn Phe Asn Ser Glu Thr
      165              170              175

Val Pro Gln Lys Ser Ser Leu Glu Glu Pro Asp Phe Tyr Lys Thr Lys
      180              185              190

Ile Lys Leu Cys Ile Leu Leu His Ala Phe Arg Ile Arg Ala Val Thr
      195              200              205

Ile Asp Arg Val Thr Ser Tyr Leu Asn Ala Ser
      210              215

```

<210> 19

<211> 132

<212> PRT

<213> Homo sapiens

<400> 19

```

Met Ala Leu Leu Leu Thr Thr Val Ile Ala Leu Thr Cys Leu Gly Gly
 1              5              10              15

Phe Ala Ser Pro Gly Pro Val Pro Pro Ser Thr Ala Leu Arg Glu Leu
      20              25              30

Ile Glu Glu Leu Val Asn Ile Thr Gln Asn Gln Lys Ala Pro Leu Cys
      35              40              45

Asn Gly Ser Met Val Trp Ser Ile Asn Leu Thr Ala Gly Met Tyr Cys
      50              55              60

```

Ala Ala Leu Glu Ser Leu Ile Asn Val Ser Gly Cys Ser Ala Ile Glu
 65 70 75 80

Lys Thr Gln Arg Met Leu Ser Gly Phe Cys Pro His Lys Val Ser Ala
 85 90 95

Gly Gln Phe Ser Ser Leu His Val Arg Asp Thr Lys Ile Glu Val Ala
 100 105 110

Gln Phe Val Lys Asp Leu Leu Leu His Leu Lys Lys Leu Phe Arg Glu
 115 120 125

Gly Arg Phe Asn
 130

<210> 20
 <211> 114
 <212> PRT
 <213> Homo sapiens

<400> 20
 Asn Trp Val Asn Val Ile Ser Asp Leu Lys Lys Ile Glu Asp Leu Ile
 1 5 10 15

Gln Ser Met His Ile Asp Ala Thr Leu Tyr Thr Glu Ser Asp Val His
 20 25 30

Pro Ser Cys Lys Val Thr Ala Met Lys Cys Phe Leu Leu Glu Leu Gln
 35 40 45

Val Ile Ser Leu Glu Ser Gly Asp Ala Ser Ile His Asp Thr Val Glu
 50 55 60

Asn Leu Ile Ile Leu Ala Asn Asn Ser Leu Ser Ser Asn Gly Asn Val
 65 70 75 80

Thr Glu Ser Gly Cys Lys Glu Cys Glu Glu Leu Glu Glu Lys Asn Ile
 85 90 95

Lys Glu Phe Leu Gln Ser Phe Val His Ile Val Gln Met Phe Ile Asn
 100 105 110

Thr Ser

<210> 21
 <211> 252
 <212> PRT
 <213> Homo sapiens

<400> 21
 Met Gly Val Leu Leu Thr Gln Arg Thr Leu Leu Ser Leu Val Leu Ala
 1 5 10 15

Leu Leu Phe Pro Ser Met Ala Ser Met Ala Ala Ile Gly Ser Cys Ser
 20 25 30

Lys Glu Tyr Arg Val Leu Leu Gly Gln Leu Gln Lys Gln Thr Asp Leu
 35 40 45
 Met Gln Asp Thr Ser Arg Leu Leu Asp Pro Tyr Ile Arg Ile Gln Gly
 50 55 60
 Leu Asp Val Pro Lys Leu Arg Glu His Cys Arg Glu Arg Pro Gly Ala
 65 70 75 80
 Phe Pro Ser Glu Glu Thr Leu Arg Gly Leu Gly Arg Arg Gly Phe Leu
 85 90 95
 Gln Thr Leu Asn Ala Thr Leu Gly Cys Val Leu His Arg Leu Ala Asp
 100 105 110
 Leu Glu Gln Arg Leu Pro Lys Ala Gln Asp Leu Glu Arg Ser Gly Leu
 115 120 125
 Asn Ile Glu Asp Leu Glu Lys Leu Gln Met Ala Arg Pro Asn Ile Leu
 130 135 140
 Gly Leu Arg Asn Asn Ile Tyr Cys Met Ala Gln Leu Leu Asp Asn Ser
 145 150 155 160
 Asp Thr Ala Glu Pro Thr Lys Ala Gly Arg Gly Ala Ser Gln Pro Pro
 165 170 175
 Thr Pro Thr Pro Ala Ser Asp Ala Phe Gln Arg Lys Leu Glu Gly Cys
 180 185 190
 Arg Phe Leu His Gly Tyr His Arg Phe Met His Ser Val Gly Arg Val
 195 200 205
 Phe Ser Lys Trp Gly Glu Ser Pro Asn Arg Ser Arg Arg His Ser Pro
 210 215 220
 His Gln Ala Leu Arg Lys Gly Val Arg Arg Thr Arg Pro Ser Arg Lys
 225 230 235 240
 Gly Lys Arg Leu Met Thr Arg Gly Gln Leu Pro Arg
 245 250

<210> 22

<211> 200

<212> PRT

<213> Homo sapiens

<400> 22

Met Ala Phe Thr Glu His Ser Pro Leu Thr Pro His Arg Arg Asp Leu
 1 5 10 15
 Cys Ser Arg Ser Ile Trp Leu Ala Arg Lys Ile Arg Ser Asp Leu Thr
 20 25 30
 Ala Leu Thr Glu Ser Tyr Val Lys His Gln Gly Leu Asn Lys Asn Ile
 35 40 45
 Asn Leu Asp Ser Ala Asp Gly Met Pro Val Ala Ser Thr Asp Gln Trp
 50 55 60

Ser Glu Leu Thr Glu Ala Glu Arg Leu Gln Glu Asn Leu Gln Ala Tyr
 65 70 75 80
 Arg Thr Phe His Val Leu Leu Ala Arg Leu Leu Glu Asp Gln Gln Val
 85 90 95
 His Phe Thr Pro Thr Glu Gly Asp Phe His Gln Ala Ile His Thr Leu
 100 105 110
 Leu Leu Gln Val Ala Ala Phe Ala Tyr Gln Ile Glu Glu Leu Met Ile
 115 120 125
 Leu Leu Glu Tyr Lys Ile Pro Arg Asn Glu Ala Asp Gly Met Pro Ile
 130 135 140
 Asn Val Gly Asp Gly Gly Leu Phe Glu Lys Lys Leu Trp Gly Leu Lys
 145 150 155 160
 Val Leu Gln Glu Leu Ser Gln Trp Thr Val Arg Ser Ile His Asp Leu
 165 170 175
 Arg Phe Ile Ser Ser His Gln Thr Gly Ile Pro Ala Arg Gly Ser His
 180 185 190
 Tyr Ile Ala Asn Asn Lys Lys Met
 195 200

<210> 23
 <211> 181
 <212> PRT
 <213> Homo sapiens

<400> 23
 Ser Pro Leu Pro Ile Thr Pro Val Asn Ala Thr Cys Ala Ile Arg His
 1 5 10 15
 Pro Cys His Asn Asn Leu Met Asn Gln Ile Arg Ser Gln Leu Ala Gln
 20 25 30
 Leu Asn Gly Ser Ala Asn Ala Leu Phe Ile Leu Tyr Tyr Thr Ala Gln
 35 40 45
 Gly Glu Pro Phe Pro Asn Asn Leu Asp Lys Leu Cys Gly Pro Asn Val
 50 55 60
 Thr Asp Phe Pro Pro Phe His Ala Asn Gly Thr Glu Lys Ala Lys Leu
 65 70 75 80
 Val Glu Leu Tyr Arg Ile Val Val Tyr Leu Gly Thr Ser Leu Gly Asn
 85 90 95
 Ile Thr Arg Asp Gln Lys Ile Leu Asn Pro Ser Ala Leu Ser Leu His
 100 105 110
 Ser Lys Leu Asn Ala Thr Ala Asp Ile Leu Arg Gly Leu Leu Ser Asn
 115 120 125
 Val Leu Cys Arg Leu Cys Ser Lys Tyr His Val Gly His Val Asp Val

130 135 140
 Thr Tyr Gly Pro Pro Asp Thr Ser Gly Lys Asp Val Phe Gln Lys Lys
 145 150 155 160
 Lys Leu Gly Cys Gln Leu Leu Gly Lys Tyr Lys Gln Ile Ile Ala Val
 165 170 175
 Leu Ala Gln Ala Phe
 180

<210> 24
 <211> 39
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:PCR Primer

<400> 24
 gggggtcgac catatgttcc caaccattcc cttatccag 39

<210> 25
 <211> 33
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:PCR Primer

<400> 25
 gggggatcct cactagaagc cacagctgcc etc 33

<210> 26
 <211> 39
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:PCR Primer

<400> 26
 ccccgatcc gccaccatgg atctctggca gctgctgtt 39

<210> 27
 <211> 40
 <212> DNA
 <213> Artificial Sequence

<220>
 <223> Description of Artificial Sequence:PCR Primer

<400> 27
 ccccgtcgac tctagagcta ttaaatacgt agctcttggg 40

<210> 28
 <211> 32
 <212> DNA
 <213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:PCR Primer

<400> 28

cgcggatccg attagaatcc acagctcccc tc

32

<210> 29

<211> 66

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:PCR Primer

<400> 29

ccccctctag acatatgaag aagaacatcg cattcctgct ggcattctatg ttctgtttct 60
ctatcg 66

<210> 30

<211> 65

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:PCR Primer

<400> 30

gcattctatgt tcgttttctc tatcgctacc aacgcttacg cattcccaac cattccctta 60
tccag 65

<210> 31

<211> 62

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:PCR Primer

<400> 31

gcagtggcac tggctggttt cgctaccgta ggcgaggcct tcccaaccat tcccttatcc 60
ag 62

<210> 32

<211> 59

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:PCR Primer

<400> 32

ccccgtcgac acatatgaag aagacagcta tcgctgattgc agtggcactg gctgggttc 59

<210> 33

<211> 36

<212> DNA

<213> Artificial Sequence

<220>

<223> Description of Artificial Sequence:PCR Primer

<400> 33
ctgcttgaag atctgcccac accggggggt gccatc 36

<210> 34
<211> 24
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:PCR Primer

<400> 34
gtagcgcagg ccttcccaac catt 24

<210> 35
<211> 39
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:PCR Primer

<400> 35
ctgcttgaag atctgcccag tccgggggca gccatcttc 39

<210> 36
<211> 51
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:PCR Primer

<400> 36
gggcagatct tcaagcagac ctacagcaag ttcgactgca actcacacaa c 51

<210> 37
<211> 34
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:PCR Primer

<400> 37
cgcggtaccc gggatccgat tagaatccac agct 34

<210> 38
<211> 36
<212> DNA
<213> Artificial Sequence

<220>
<223> Description of Artificial Sequence:PCR Primer

<400> 38
gggcagatct tcaagcagac ctactgcaag ttcgac 36

<210> 39
<211> 42
<212> DNA

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US98/14497

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) :Please See Extra Sheet.

US CL :Please See Extra Sheet.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 530/351; 435/69.1, 69.5, 69.51, 69.52, 71.1, 71.2, 325, 252.3, 320.1

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched
NONE

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

APS, CAS ONLINE, MEDLINE, CAPLUS, BIOSIS

search terms: cysteine variant, derivative, analog, growth hormone, substitution, glycosylation

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 4,588,585 A (MARK et al.) 13 May 1986, see entire document.	1, 3, 6, 9-11
A	US 5,208,158 A (BECH et al.) 04 May 1993, see entire document.	1, 3, 6, 9-11
A	US 5,166,322 A (SHAW et al.) 24 November 1992, see entire document.	1, 3, 6, 9-11
A	US 5,206,344 A (KATRE et al.) 27 April 1993, see entire document.	1, 3, 6, 9-11

☐ Further documents are listed in the continuation of Box C.
 ☐ See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
B earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*G* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

13 OCTOBER 1998

Date of mailing of the international search report

22 OCT 1998

 Name and mailing address of the ISA/US
 Commissioner of Patents and Trademarks
 Box PCT
 Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

PREMA MERTZ

Telephone No. (703) 308-0196

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US98/14497**Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)**

This international report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☒ Claims Nos.: 5, 8 and 23
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

Please See Extra Sheet.

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:
1, 3, 6, 9-11

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US98/14497

A. CLASSIFICATION OF SUBJECT MATTER: IPC (6):

C07K 14/475, 14/52, 14/53, 14/535, 14/54, 14/55, 14/555, 14/56, 14/565, 14/57; C12N 15/09, 15/12, 15/18, 15/19, 15/20, 15/21, 15/22, 15/23, 15/24, 15/26, 15/27

A. CLASSIFICATION OF SUBJECT MATTER: US CL :

530/351; 435/69.1, 69.5, 69.51, 69.52, 71.1, 71.2, 325, 252.3, 320.1

BOX II. OBSERVATIONS WHERE UNITY OF INVENTION WAS LACKING

This ISA found multiple inventions as follows:

This application contains the following inventions or groups of inventions which are not so linked as to form a single inventive concept under PCT Rule 13.1.

Group I, claims 1, 3, 6, 9-11, drawn to cysteine variants comprising cysteine residue substitutions of growth hormone.
Group II, claims 1, 3, 6, 12-14, drawn to cysteine variants comprising cysteine residue substitutions of erythropoietin.
Group III, claims 2, 4, 7, 17-18, drawn to cysteine variants comprising cysteine residue insertions of erythropoietin.
Group IV, claims 2, 4, 7, 15-16, drawn to cysteine variants comprising cysteine residue insertions of growth hormone.
Group V, claims 2, 4, 7, 19-20, drawn to cysteine variants comprising cysteine residue insertions of alpha interferon.
Group VI, claims 2, 4, 7, 21-22, drawn to cysteine variants comprising cysteine residue insertions of granulocyte-colony stimulating factor.

The inventions listed as Groups I-VI do not relate to a single inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons:

The inventions listed as Groups I-VI do not relate to a single inventive concept under PCT Rule 13.1 because, under PCT Rule 13.2, they lack the same or corresponding special technical features for the following reasons:

Pursuant to 37 C.F.R. § 1.475 (d), the ISA/US considers that where multiple products and processes are claimed, the main invention shall consist of the first invention of the category first mentioned in the claims and the first recited invention of each of the other categories related thereto. Accordingly, the main invention (Group I) comprises the first-recited product, a cysteine variant comprising cysteine residue substitutions in growth hormone. Further pursuant to 37 C.F.R. § 1.475 (d), the ISA/US considers that any feature which the subsequently recited products and methods share with the main invention does not constitute a special technical feature within the meaning of PCT Rule 13.2 and that each of such products and methods accordingly defines a separate invention.